

Statistics of X-ray Observables of Cooling Core and Non-cooling Core Clusters of Galaxies

Y. Chen^{1,2}, T.H. Reiprich³, H. Böhringer², Y. Ikebe⁴, and Y.-Y. Zhang²

¹ Key Laboratory of Particle Astrophysics, Institute of High Energy Physics, Chinese Academy of Sciences, Beijing 100049, P.R. China

² Max-Planck-Institut für Extraterrestrische Physik, D-85748 Garching, Germany

³ Argelander-Institut für Astronomie, Universität Bonn, Auf dem Hügel 71, 53121 Bonn, Germany

⁴ National Museum of Emerging Science and Innovation, Tokyo, 135-0064, Japan

Received ?; accepted ?

Abstract. We present a statistical study of the occurrence and effects of the cooling cores in clusters of galaxies in a flux limited sample, HIFLUGCS, based on ROSAT and ASCA observations. About 49% of the clusters in this sample have a significant, classically calculated cooling flow mass deposition rate. The upper envelope of the derived mass deposition rate is roughly proportional to cluster mass, and the fraction of cooling core clusters is found to decrease with the cluster mass. The cooling core clusters are found to have smaller core radii than non-cooling core clusters, while some non-cooling core clusters have large β values (> 0.8). In the relation of the X-ray luminosity v.s. the temperature and the mass, the cooling core clusters show a significantly higher normalization. A systematic correlation analysis, involving also relations of the gas mass and the total infrared luminosity, indicates that this bias is shown to be mostly due to an enhanced X-ray luminosity of cooling core clusters, while the other parameters, like temperature, mass, and gas mass may be less affected by the occurrence of a cooling core. These results may be explained by the fact that at least some of the non-cooling core clusters are in dynamically young states compared with cooling core clusters and they may turn into cooling core clusters in a later evolutionary stage.

Key words. Galaxies: clusters: general – Galaxies: clusters: Intergalactic medium – X-ray: galaxies: clusters

1. Introduction

Clusters of galaxies are interesting large-scale astrophysical laboratories and ideal probes for the study of the large-scale structure of the Universe and for the test of cosmological models (e.g. Voit 2005). A very important scaling parameter that is crucial in these studies is the cluster mass, which cannot easily be measured unless detailed observations are available. It is therefore estimated by means of other suitable, easily obtained global observables like for example X-ray luminosity or X-ray temperature (e.g. Reiprich & Böhringer 2002, Markevitch 1998, Ikebe et al. 2002, Finoguenov et al. 2001, Arnaud et al. 2005). Since the early days of X-ray imaging with the EINSTEIN satellite it is apparent that there may be two, to some extent, distinct classes of galaxy clusters: clusters with very dense gaseous core region, so-called cooling cores, and another type with shallower cores often exhibiting more internal structure (e.g. Jones & Forman 1984, Ota & Mitsuda 2004, Peres et al. 1998, Schuecker et al. 2001a). In the present

paper we explore the influence of this dichotomy on the scaling relations between global cluster X-ray observables, and between the observables and the cluster mass in order to improve our understanding of the use of such scaling relations in cosmological applications.

Clusters with dense gaseous cores, with central cooling times significantly lower than a Hubble time, have formerly been termed cooling flow clusters, and it was believed that the intracluster medium (ICM) in these regions cools and condenses, as it is difficult to avoid cooling in the absence of a very fine-tuned heating mechanism (Fabian 1994). A different point of view, not requiring a cooling flow, has been put forward early based on ASCA spectroscopic results (e.g., Ikebe et al. 1999; Makishima et al. 2001, and references therein). With XMM-Newton observations it was found that the spectral features predicted by the classical cooling flow model are not observed in the X-ray spectra of cooling flow regions (e.g. Peterson et al. 2001, 2003). While a slight temperature decrease by factors up to 2 - 3 towards the center in cooling cores is observed, the expected features for further cooling are absent. With high resolution Chandra observations a pos-

sible fine-tuned heat source has been found in the interaction of central AGN with the cluster ICM, which is now taken as the most probable reason for the prevention of massive cooling flows (e.g. David et al. 2001, Böhringer et al. 2002, Fabian et al. 2003, McNamara et al. 2005). Therefore we follow the now widely used convention to call the clusters in our sample cooling core clusters (CCC) and non-cooling core clusters (NCCC).

An influence of the CCC or NCCC nature of the clusters on the scaling relations of global properties has previously been realized, e.g. in the luminosity temperature relation (e.g. Fabian et al. 1994, Markevitch 1998, McCarthy et al. 2004) and other parameters (O’Hara et al. 2006). Here we extend the studies of the influence of CCCs on the scaling relation to the largest X-ray flux limited sample of galaxy cluster with detailed X-ray data that allow a mass determination, the HIFLUGCS (the HIghest X-ray FLUX Galaxy Cluster Sample; Reiprich 2001, Reiprich & Böhringer 2002). This cluster sample is selected only by X-ray flux irrespective of the cluster morphology and we do not know of any signature of incompleteness of the sample. Therefore it should provide a representative mix of cluster morphologies for given X-ray luminosity providing the correct statistics to be applied to the typical cosmological X-ray survey cluster samples.

In particular we study the segregation of CC and NCC clusters in the $L_X - T$, $L_X - M$, $M - T$, and the $f_{gas} - T$ relation. A major goal in this study is to better understand the scatter in these relations which has to be folded into the test of large-scale structure measures and cosmological models (e.g., Ikebe et al. 2002; Stanek et al. 2006). It is especially interesting in the context of the $L_X - M$ relation given by Reiprich & Böhringer (2002) where the observed scatter is very large and partly due to the large mass determination uncertainties. Therefore it was very difficult to separate the intrinsic scatter from the scatter introduced by the formal and systematic measurement errors. It is the intrinsic scatter, however, that is important for the application. A difference in the relation amplitude between CCCs and NCCCs could in principle provide a lower limit to the intrinsic scatter in the $L_X - M$ relation of Reiprich & Böhringer (2002) if the systematic uncertainties are well controlled and thus help to further understand the origin of the scatter. This is interesting because the best fit cosmological parameter values from the WMAP 3rd year data (Spergel et al. 2006) applied to compare the predicted and observational derived HIFLUGCS cluster mass function provide an indication that the intrinsic scatter is probably smaller than the systematic measurement errors that go into the derived mass and X-ray luminosity relation (Reiprich 2006). Our study has a lot in common with the work of O’Hara et al. (2006), but was started independently a few years ago, and we discuss the correspondence of the two studies throughout the paper.

The paper is organized as follows: In Sect.2 we briefly introduce the sample. In Sect.3, we present the method of data reduction. We compare the properties of the CCC and NCCC in Sect.4 and discuss the implications of the

results. Sect.5 provides a summary. In the following we adopt a cosmological model with $H_0 = 50 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_m=1$ and $\Omega_\Lambda=0$, a choice which was mostly made for easier comparison with previous results.

2. The sample

The extended HIFLUGCS sample with 106 clusters and groups of galaxies is used for the present study. 92 of these clusters have known temperatures determined from X-ray spectroscopy. Here we use two cluster temperatures: T_m , the emission measure weighted temperature, which is mainly derived from a single temperature fit to the global X-ray spectrum of the clusters (Markevitch et al. 1998, Reiprich 2001 and references therein), and T_h , which is the hotter bulk component of a two-temperature model fitted to the spectrum (88 of them from ASCA, Ikebe et al. 2002). T_h was determined by accepting a small second lower temperature component, to allow for a low temperature phase in a possible cooling core in the central cluster region. The typical temperature of this second component was about a factor of 2 lower than the bulk temperature (Ikebe et al. 2002). The second component of the two-temperature fit, which is generally only needed for the cooling core clusters, has a small normalization and is expected to account for the lower central temperature phase in the cooling cores. For the clusters with no measured T_m or T_h , we derived them using the $L_X - T$ relation of Markevitch (1998) with $L_X (< 2 \text{ Mpc})$ uncorrected and T corrected for cooling flows. The hotter component, T_h , which is usually slightly higher than T_m , is expected to provide a good measure of the gravitational potential depth and the total mass of the clusters.

The X-ray surface brightness profiles are derived from ROSAT PSPC observations, 36 of them are from RASS observations (allowing for a large enough field-of-view for the prominent nearby clusters) and 70 from pointing observations (Reiprich & Böhringer 2002). The large FOV of the ROSAT PSPC allows us to cover most of the clusters out to r_{500} , the radius at which the mean density of the cluster is 500 times of that of a critical density universe. In addition, it is worth to note that this flux limited sample has the largest sky area so far.

The basic properties of the clusters in the HIFLUGCS sample from Reiprich & Böhringer (2002) are given in Table 1 and 2. Note that T_m , the emission measure weighted temperature, is used in the calculation of some quantities such as n_{center} , t_{cool} and \dot{M} in the next section. In Fig. 1 we show the distribution of the masses and cooling flow mass deposition rates determined as described in the following sections. The NCC clusters with no significant mass deposition rates and very small or no cooling radii are shown with their mass distribution at the bottom of the plot. A striking feature of the plot is the ridge of cooling core clusters with formal mass deposition rates that increase almost linearly with cluster mass. This leads us to define the class of pronounced cooling core clusters by a lower limit of the ratio of the formal mass deposition




Fig. 1. Formally deduced mass deposition rates in the frame of the classical cooling flow model as a function of the total cluster mass, M_{500} . There is a pronounced ridge line of stronger cooling core clusters with a formal mass deposition rate almost proportional to the cluster mass. The clusters with insignificant mass deposition rates below a value of $0.01 M_{\odot} \text{ yr}^{-1}$ are plotted according to their total mass at the bottom of the plot at a formal value of $0.01 M_{\odot} \text{ yr}^{-1}$. The line at $\dot{M}/M_{500} = 10^{-13} \text{ yr}^{-1}$ separates strong CCCs from small to moderate CCCs. In all figures of this paper, the filled black circles represent the pronounced CCCs. The filled grey circles are small to moderate CCCs and the open circles are NCCCs. Throughout this paper the error bars are at $\pm 68\%$ confidence level except for T_m of some clusters, T_h and L_X which are at 90% confidence level (and thus a conservative error estimate).

rate to the cluster mass, M_{500} . The ratio value chosen of \dot{M}/M_{500} of 10^{-13} yr^{-1} is indicated in Fig. 1. A further limit in the mass deposition rate at $0.01 M_{\odot} \text{ yr}^{-1}$ is used to separate small to moderate cooling cores from NCCC. The total sample thus splits up in 36 pronounced CCCs, 16 small to moderate CCCs, and 54 NCCCs. We will make use of this classification in the further discussion.

3. Data analysis

3.1. X-ray surface brightness and mass profiles

To determine the X-ray surface brightness distribution we produced images in the 0.5 to 2 keV band (PSPC PI channel 52 to 201) and the corresponding vignetting corrected exposure maps. Contaminating point sources and obvious substructure were excised. A center position was obtained from an iterative determination of the “center-of-mass” of the photon distribution in a 7.5 arcmin radius aperture. The surface brightness profile was then constructed by azimuthal averaging in concentric bins. This procedure is the same as used in Reiprich & Böhringer (2002).

We fit the surface brightness profile with a single β model (Cavaliere & Fusco-Femiano 1976)

$$S(r) = S_0 \left(1 + (r/r_c)^2\right)^{-3\beta+1/2}, \quad (1)$$

where S_0 is central brightness (counts/s/pixel²; $1' = 120$ pixels for ROSAT PSPC) and r_c the core radius (kpc). We also try a double β model

$$S(r) = S_{01}\alpha_1^{-3\beta_1+1/2} + S_{02}\alpha_2^{-3\beta_2+1/2}, \quad (2)$$

$$\alpha_1 = 1 + (r/r_{c1})^2, \quad (3)$$

$$\alpha_2 = 1 + (r/r_{c2})^2, \quad (4)$$

in which the fits of 49 clusters have significantly improved reduced χ^2 values compared to the fits using a single β model (Table 3). We thus use the double β model for these clusters. Assuming that the temperature is homogeneous in the cluster, we can calculate the gas number density profile $n(r)$. The errors introduced by this simplification in the presence of temperature variations is only of the order a few percent which justifies this approximation. Assuming moreover that the gas is in hydrostatic equilibrium, the total mass of the cluster can be calculated for a single β model as

$$M(r) = \frac{3\beta k T_h r}{G\mu m_p} \frac{(r/r_c)^2}{1 + (r/r_c)^2}, \quad (5)$$

where k is the Boltzmann constant, G the gravitational constant, μ the molecular weight ($\mu=0.61$), and m_p the proton mass. For a double β model we find

$$M(r) = \frac{3kT_h r^3}{G\mu m_p} \frac{n_{01}^2 \beta_1 \alpha_1^{-3\beta_1-1}/r_{c1}^2 + n_{02}^2 \beta_2 \alpha_2^{-3\beta_2-1}/r_{c2}^2}{n_{01}^2 \alpha_1^{-3\beta_1} + n_{02}^2 \alpha_2^{-3\beta_2}}, \quad (6)$$

where n_{01} and n_{02} is the central equivalent electron number density calculated from the two surface brightness components, respectively. The central electron number density, n_0 , can be derived from

$$n_0^2 = n_{01}^2 + n_{02}^2. \quad (7)$$

3.2. Cooling core properties

The cooling time of the gas is calculated by

$$t_{cool} = \frac{5}{2} \frac{n_e + n_i}{n_e} \frac{kT_m}{n_H \Lambda(A, T_m)}, \quad (8)$$

where $\Lambda(A, T_m)$ is the cooling function of the gas, n_e , n_i and n_H are the number densities of the electrons, ions, and hydrogen, respectively. Here we use the abundance $A = 0.3$ for all clusters. Note that for the nearly fully ionized plasma in clusters, $n_e = 1.2n_H$ and $n_i = 1.1n_H$.

Following the previously most frequently used convention, we define the cooling radius as the radius where the gas cooling time is equal to the age of the cluster assumed to be close to the Hubble time ($t_{age} \sim 1/H_0 = 13 \text{ Gyr}$). The physical meaning of the cooling radius within the classical cooling flow model is, that within the cooling radius the gas will lose all of its energy by X-ray emission and is replaced by ambient hot gas from larger radii in a steady state inflow. We can therefore calculate the energy loss rate from the integral of the X-ray emission inside the

cooling radius and the mass inflow rate from the enthalpy influx necessary to compensate for this energy loss. We further account for the energy gain as the inflowing gas moves down the gravitational potential gradient. Thus, the total mass deposition rate within the shell i can be determined by

$$\dot{M}(i) = \frac{n_e(i)n_H(i)\Lambda(A, T_m)V(i) + \frac{5}{2}\frac{kT_m}{\mu m_p}\dot{M}(i-1)}{\frac{5}{2}\frac{kT_m}{\mu m_p} + \Phi(i+1) - \Phi(i)}, \quad (9)$$

where $V(i)$ is volume of the shell i , $\Phi(i+1)$ and $\Phi(i)$ is the gravitational potential of the shell $i+1$ and i respectively. For a single β model, $\Phi(i)$ can be calculated from

$$\Phi(i) = \frac{3}{2}\frac{\beta kT_h}{\mu m_p} \ln\left(1 + \left(\frac{r(i)}{r_c}\right)^2\right). \quad (10)$$

When we use the double β model, these formulae will change accordingly, and we spare their writing-out for brevity.

3.3. Error estimate

We adopted a Monte-Carlo method to estimate the errors of the cluster properties derived in the previous subsection, such as the mass and mass deposition rate. In the calculation of the mass of the clusters, we assumed a polytropic index with a value of $\gamma = 1$. From previous observational studies the range of γ -values is constrained to be between 0.9 and 1.3 (e.g. Finoguenov 2001, Pratt et al. 2006). The main errors are from the temperature and the γ . We use a $\beta\gamma$ model (e.g. Ettori 2000) to estimate the errors of the mass deposition rate and we assume a polytropic index greater than 0.9 and following a Gaussian distribution with a mean value of 1.15 and a variance of 0.15 as derived in Finoguenov et al. (2001). For each cluster in the HIFLUGCS, we created a cluster sample with 1000 clusters with simulated T_m , T_h , S , β , r_c and γ according to their own errors. We calculated other properties (e.g. n , t_{cool} , \dot{M} , M and so on) of each simulated cluster and then obtained the errors.

4. Statistical properties

4.1. Basic Properties

In the following we investigate the relations between several observables and the cluster mass. For all the relations we use the BCES-Bisector fit of Akritas & Bershady (1996). The fits are performed to the logarithmic values of the parameters and quoted in the form

$$\log_{10}(Y) = A + B \cdot \log_{10}(X)$$

in Table 7. The median X-ray luminosity, $L_X(0.1 - 2.4 \text{ keV})$, the median temperature, T_m , and the median cluster mass, M_{500} of the sample are $2.9 \cdot 10^{44} \text{ erg s}^{-1}$, 4.1 keV, and $4.8 \cdot 10^{14} M_\odot$, respectively. Therefore we use the values $1 \cdot 10^{44} \text{ erg s}^{-1}$, 4 keV, and $5 \cdot 10^{14} M_\odot$, respectively, as pivot points for the fits of the relations.




Fig. 2. The numbers of CCCs (solid lines) and NCCCs (dashed lines) versus the mass of the clusters M_{500} . Note that the fraction of NCCCs increases with M_{500} .

In Fig. 2 we show the number of CCCs (here including the strong CCC and moderate CCC) and NCCCs in the sample as a function of the cluster mass. The fraction of NCCCs clearly increases with M_{500} . This is also seen in the smaller flux limited sample used by O'Hara et al. (2006). At the low M_{500} end, this may partly be due to the fact that some small mass non-cooling core groups may have small luminosities and do not reach the flux limit of HIFLUGCS. In general, however, the main reasons may be that the fraction of dynamically young clusters increases with cluster mass and these clusters in general do not feature cooling cores. In addition the ICM is hotter in more massive clusters making the radiative heat loss relatively slower. This is an important statistical property of the cluster sample to keep in mind, since any segregation of CCCs and NCCCs in the parameter relations can then also introduce a mass dependent effect in the relations of the combined sample.

Fig. 3 shows the distribution of the values for the core radius, r_c , and slope parameter β from the fit of the β -model to the X-ray surface brightness profiles of the clusters in the sample. Similar to previous work (e.g. Jones & Forman 1984, White et al. 2000, Ota & Mitsuda 2004), the CC clusters segregate very distinctly at smaller values of the core radius than the NCC clusters. Furthermore, we find that the larger the relative \dot{M}_r , the smaller the r_c . In addition, it was found that some NCCCs have large β values with $\beta > 0.8$ while such high values are not found among the CCCs. Two reasons explain this behavior. For similar gravitational potential shapes in CCCs and NCCCs, the CC clusters with central temperature drops and a corresponding central ICM density increase in pressure equilibrium feature X-ray surface brightness cusps which are fit by smaller core radii (Jones & Forman 1984). In addition NCCCs are often dynamically young featuring substructure, elongations, or disturbed core regions which result in inflated core radii, which in turn lead to steeper outer surface brightness slopes.




Fig. 3. r_c - β diagram. Note that the cooling-core clusters have smaller core radii.




Fig. 4. r_c - M_{500} diagram. r_c increases with M_{500} faster than the self-similar expectation $r_c \propto M_{500}^{1/3}$. The solid line represents the BCES bisector fit for all 106 clusters with the form $\log_{10} \left(\frac{r_c}{100 \text{ kpc}} \right) = A + B \cdot \log_{10} \left(\frac{M_{500}}{510^{14} M_{\odot}} \right)$. The dashed line and the dotted line represent the fits for the NCCCs and CCCs, respectively (see Table 6).

In Fig. 4 we explore the relation of the core radius, r_c , with the cluster mass, M_{500} . Assuming that clusters have a strictly self-similar shape, we expect that any characteristic radius scales as $r \propto M^{1/3}$. The results of power law scaling relation fits to the data are given in Table 6 and are shown in Fig. 4. For both subsamples, CCCs and NCCCs, the observed slope is steeper than this simple expectation, that is the core radius increases faster with mass than expected. The explanation for this behavior is probably not trivial. For the NCCCs the reason might be again that the fraction of dynamically young clusters with inflated core radii may be larger for higher cluster masses. For the CCCs it might be the increasing dominance of the central cluster galaxy with decreasing cluster mass that makes the core region relatively more compact for less massive systems.




Fig. 5. M_{500} - T_h relation. The slopes derived from the total 88 clusters with measured T_h from ASCA (solid line), CCCs (dotted line) and NCCCs (dashed line) are consistent with each other.

It is interesting to note that the relation fitted to the complete sample is steeper than each of the separately fitted relations. This is exactly the effect mentioned above. It is the result of having an offset in the relation of the two subsamples (significantly smaller core radii for the CCCs) and in addition a biased distribution of the clusters in the two subsamples with more CCCs at the low mass end and more NCCCs at the high mass end. Among all the plots we show in this paper, this is the relation where this effect is most pronounced.

4.2. M - T_h relation

Fig. 5 shows the relation of the globally measured X-ray temperature of the hot component, T_h (excluding non-ASCA derived T_h), and the cluster mass, M_{500} . The self-similar model prediction is a slope with a value of 1.5, consistent with the value we obtain for the total sample with measured T_h and consistent with the discussion in Finoguenov et al. (2001). The results of power law fits to all the relations discussed in this section are summarized in Table 7. Moreover, the slopes and the normalizations of the M - T_h relation for the CCCs and NCCCs are consistent within errors. Thus there is no significant influence of cooling cores on this relation. Note, however, that the two cluster parameters compared in this relation, T_h and M_{500} are not independently obtained, but M_{500} is directly dependent on the temperature measurements. If T_h has an offset ΔT_h , M_{500} will change to $M_{500} + \Delta M_{500} \propto (T_h + \Delta T_h)^{3/2}$. Note that this slope is the same as the self-similar model prediction. Thus any influence of the effects of cooling cores on the temperature determination will also appear in the cluster mass

Table 1. Cluster properties.

Name	z	T_m (keV)	β	r_c (kpc)	n_{center} (10^{-2} cm $^{-3}$)	t_{cool} (10^{10} yr)	\dot{M} (M_{\odot}/yr)
2A0335	0.0349	$3.01^{+0.07}_{-0.07}$	$0.575^{+0.004}_{-0.003}$	33^{+0}_{-0}	$5.47^{+0.06}_{-0.06}$	$0.10^{+0.00}_{-0.00}$	360^{+20}_{-17}
A0085	0.0556	$6.10^{+0.20}_{-0.20}$	$0.532^{+0.004}_{-0.004}$	82^{+3}_{-3}	$2.57^{+0.10}_{-0.10}$	$0.33^{+0.02}_{-0.02}$	200^{+33}_{-27}
A0119	0.0440	$5.80^{+0.60}_{-0.60}$	$0.675^{+0.026}_{-0.023}$	501^{+27}_{-26}	$0.15^{+0.01}_{-0.01}$	$5.49^{+0.53}_{-0.53}$	0^{+0}_{-0}
A0133	0.0569	$3.80^{+0.90}_{-0.90}$	$0.530^{+0.004}_{-0.018}$	45^{+1}_{-1}	$2.80^{+0.08}_{-0.08}$	$0.23^{+0.09}_{-0.07}$	108^{+55}_{-51}
A0262	0.0161	$2.15^{+0.06}_{-0.06}$	$0.443^{+0.018}_{-0.017}$	41^{+11}_{-9}	$0.81^{+0.13}_{-0.09}$	$0.56^{+0.04}_{-0.08}$	14^{+1}_{-4}
A0399	0.0715	$7.40^{+0.70}_{-0.70}$	$0.713^{+0.137}_{-0.095}$	449^{+131}_{-99}	$0.22^{+0.04}_{-0.03}$	$4.32^{+0.78}_{-0.70}$	0^{+0}_{-0}
A0400	0.0240	$2.31^{+0.14}_{-0.14}$	$0.534^{+0.014}_{-0.013}$	154^{+9}_{-8}	$0.20^{+0.01}_{-0.01}$	$2.39^{+0.13}_{-0.12}$	0^{+0}_{-0}
A0401	0.0748	$8.30^{+0.50}_{-0.50}$	$0.613^{+0.010}_{-0.010}$	245^{+11}_{-10}	$0.60^{+0.05}_{-0.04}$	$1.67^{+0.15}_{-0.14}$	0^{+0}_{-0}
A0478	0.0900	$7.10^{+0.40}_{-0.40}$	$0.613^{+0.004}_{-0.004}$	98^{+2}_{-2}	$3.55^{+0.15}_{-0.14}$	$0.26^{+0.02}_{-0.01}$	645^{+137}_{-113}
A0496	0.0328	$4.13^{+0.08}_{-0.08}$	$0.484^{+0.003}_{-0.003}$	30^{+1}_{-1}	$4.07^{+0.29}_{-0.29}$	$0.17^{+0.01}_{-0.01}$	114^{+35}_{-28}
A0576	0.0381	$4.02^{+0.07}_{-0.07}$	$0.825^{+0.432}_{-0.185}$	394^{+221}_{-125}	$0.16^{+0.06}_{-0.04}$	$4.27^{+1.22}_{-1.09}$	0^{+0}_{-0}
A0754	0.0528	$9.00^{+0.50}_{-0.50}$	$0.698^{+0.027}_{-0.024}$	239^{+17}_{-16}	$0.44^{+0.02}_{-0.02}$	$2.41^{+0.13}_{-0.13}$	0^{+0}_{-0}
A1060	0.0114	$3.24^{+0.06}_{-0.06}$	$0.607^{+0.040}_{-0.034}$	94^{+15}_{-12}	$0.47^{+0.04}_{-0.04}$	$1.29^{+0.11}_{-0.11}$	0^{+0}_{-0}
A1367	0.0216	$3.55^{+0.08}_{-0.08}$	$0.695^{+0.035}_{-0.032}$	383^{+24}_{-22}	$0.15^{+0.01}_{-0.01}$	$4.29^{+0.22}_{-0.21}$	0^{+0}_{-0}
A1644	0.0474	$4.70^{+0.90}_{-0.70}$	$0.579^{+0.111}_{-0.074}$	299^{+127}_{-92}	$0.28^{+0.08}_{-0.06}$	$2.65^{+0.96}_{-0.67}$	0^{+0}_{-0}
A1650	0.0845	$5.60^{+0.60}_{-0.60}$	$0.704^{+0.131}_{-0.081}$	281^{+104}_{-70}	$0.43^{+0.09}_{-0.06}$	$1.87^{+0.33}_{-0.36}$	0^{+0}_{-0}
A1651	0.0860	$6.30^{+0.50}_{-0.50}$	$0.643^{+0.014}_{-0.013}$	180^{+9}_{-9}	$1.10^{+0.07}_{-0.07}$	$0.78^{+0.07}_{-0.07}$	81^{+30}_{-23}
A1736	0.0461	$3.50^{+0.40}_{-0.40}$	$0.542^{+0.147}_{-0.092}$	374^{+177}_{-129}	$0.13^{+0.04}_{-0.02}$	$4.92^{+1.27}_{-1.28}$	0^{+0}_{-0}
A1795	0.0616	$6.00^{+0.30}_{-0.30}$	$0.596^{+0.003}_{-0.002}$	78^{+1}_{-1}	$2.88^{+0.05}_{-0.05}$	$0.29^{+0.01}_{-0.01}$	399^{+47}_{-39}
A2029	0.0767	$8.70^{+0.30}_{-0.30}$	$0.582^{+0.004}_{-0.004}$	83^{+2}_{-2}	$3.90^{+0.15}_{-0.15}$	$0.26^{+0.01}_{-0.01}$	514^{+93}_{-78}
A2052	0.0348	$3.03^{+0.04}_{-0.04}$	$0.526^{+0.005}_{-0.005}$	36^{+1}_{-1}	$2.85^{+0.66}_{-0.62}$	$0.20^{+0.06}_{-0.04}$	108^{+188}_{-49}
A2063	0.0354	$3.68^{+0.11}_{-0.11}$	$0.561^{+0.011}_{-0.011}$	110^{+6}_{-6}	$0.96^{+0.00}_{-0.00}$	$0.67^{+0.02}_{-0.02}$	21^{+2}_{-2}
A2065	0.0721	$5.40^{+0.30}_{-0.30}$	$1.162^{+0.734}_{-0.282}$	689^{+360}_{-186}	$0.20^{+0.04}_{-0.04}$	$3.99^{+1.04}_{-1.04}$	0^{+0}_{-0}
A2142	0.0899	$8.80^{+0.60}_{-0.60}$	$0.591^{+0.006}_{-0.006}$	153^{+5}_{-5}	$1.61^{+0.07}_{-0.07}$	$0.64^{+0.04}_{-0.04}$	337^{+82}_{-61}
A2147	0.0351	$4.91^{+0.28}_{-0.28}$	$0.444^{+0.071}_{-0.046}$	237^{+102}_{-64}	$0.17^{+0.04}_{-0.03}$	$4.41^{+0.92}_{-0.87}$	0^{+0}_{-0}
A2163	0.2010	$13.29^{+0.64}_{-0.64}$	$0.796^{+0.030}_{-0.028}$	519^{+30}_{-29}	$0.53^{+0.02}_{-0.02}$	$2.37^{+0.12}_{-0.11}$	0^{+0}_{-0}
A2199	0.0302	$4.10^{+0.08}_{-0.08}$	$0.655^{+0.019}_{-0.021}$	139^{+10}_{-9}	$0.83^{+0.03}_{-0.03}$	$0.83^{+0.03}_{-0.03}$	77^{+7}_{-6}
A2204	0.1523	$7.21^{+0.25}_{-0.25}$	$0.597^{+0.008}_{-0.007}$	67^{+3}_{-2}	$5.07^{+0.12}_{-0.11}$	$0.18^{+0.01}_{-0.01}$	1287^{+122}_{-129}
A2244	0.0970	$7.10^{+5.00}_{-2.00}$	$0.607^{+0.016}_{-0.016}$	125^{+10}_{-10}	$1.20^{+0.06}_{-0.06}$	$0.76^{+0.35}_{-0.18}$	129^{+199}_{-127}
A2255	0.0800	$6.87^{+0.20}_{-0.20}$	$0.797^{+0.033}_{-0.030}$	593^{+35}_{-32}	$0.18^{+0.02}_{-0.02}$	$4.96^{+0.45}_{-0.44}$	0^{+0}_{-0}
A2256	0.0601	$7.50^{+0.40}_{-0.40}$	$0.914^{+0.054}_{-0.047}$	587^{+39}_{-36}	$0.26^{+0.01}_{-0.01}$	$3.62^{+0.19}_{-0.21}$	0^{+0}_{-0}
A2589	0.0416	$3.70^{+2.20}_{-1.10}$	$0.596^{+0.013}_{-0.012}$	118^{+7}_{-7}	$0.74^{+0.07}_{-0.07}$	$0.88^{+0.37}_{-0.24}$	19^{+53}_{-19}
A2597	0.0852	$3.60^{+0.20}_{-0.20}$	$0.633^{+0.008}_{-0.008}$	57^{+2}_{-2}	$3.63^{+0.08}_{-0.07}$	$0.17^{+0.01}_{-0.01}$	501^{+58}_{-51}
A2634	0.0312	$3.70^{+0.28}_{-0.28}$	$0.640^{+0.051}_{-0.043}$	364^{+43}_{-38}	$0.24^{+0.02}_{-0.02}$	$2.70^{+0.28}_{-0.25}$	0^{+0}_{-0}
A2657	0.0404	$3.70^{+0.30}_{-0.30}$	$0.556^{+0.007}_{-0.006}$	119^{+5}_{-4}	$0.54^{+0.00}_{-0.00}$	$1.20^{+0.07}_{-0.07}$	2^{+5}_{-5}
A3112	0.0750	$4.70^{+0.40}_{-0.40}$	$0.576^{+0.006}_{-0.006}$	61^{+2}_{-2}	$3.52^{+0.23}_{-0.23}$	$0.21^{+0.02}_{-0.02}$	346^{+123}_{-97}
A3158	0.0590	$5.77^{+0.10}_{-0.05}$	$0.661^{+0.025}_{-0.022}$	268^{+19}_{-18}	$0.39^{+0.02}_{-0.01}$	$2.11^{+0.08}_{-0.08}$	0^{+0}_{-0}
A3266	0.0594	$7.70^{+0.80}_{-0.80}$	$0.796^{+0.020}_{-0.019}$	564^{+20}_{-19}	$0.28^{+0.01}_{-0.01}$	$3.44^{+0.30}_{-0.30}$	0^{+0}_{-0}
A3376	0.0455	$4.30^{+0.60}_{-0.60}$	$1.054^{+0.101}_{-0.083}$	754^{+68}_{-60}	$0.10^{+0.01}_{-0.00}$	$6.87^{+0.83}_{-0.78}$	0^{+0}_{-0}
A3391	0.0531	$5.70^{+0.70}_{-0.70}$	$0.579^{+0.026}_{-0.024}$	234^{+23}_{-21}	$0.43^{+0.08}_{-0.06}$	$1.91^{+0.34}_{-0.34}$	0^{+0}_{-0}
A3395s	0.0498	$4.80^{+0.40}_{-0.40}$	$0.964^{+0.275}_{-0.167}$	604^{+172}_{-117}	$0.13^{+0.02}_{-0.02}$	$5.82^{+1.01}_{-0.92}$	0^{+0}_{-0}
A3526	0.0103	$3.68^{+0.06}_{-0.06}$	$0.495^{+0.011}_{-0.010}$	37^{+4}_{-4}	$1.83^{+0.13}_{-0.12}$	$0.36^{+0.03}_{-0.02}$	24^{+5}_{-5}
A3558	0.0480	$5.50^{+0.30}_{-0.30}$	$0.580^{+0.006}_{-0.005}$	223^{+5}_{-5}	$0.46^{+0.01}_{-0.01}$	$1.77^{+0.09}_{-0.09}$	0^{+0}_{-0}
A3562	0.0499	$5.16^{+0.16}_{-0.16}$	$0.472^{+0.006}_{-0.006}$	98^{+5}_{-5}	$0.58^{+0.02}_{-0.02}$	$1.33^{+0.07}_{-0.06}$	0^{+0}_{-0}
A3571	0.0397	$6.90^{+0.30}_{-0.30}$	$0.613^{+0.010}_{-0.010}$	181^{+6}_{-6}	$1.09^{+0.09}_{-0.08}$	$0.84^{+0.08}_{-0.07}$	35^{+11}_{-10}
A3581	0.0214	$1.83^{+0.04}_{-0.04}$	$0.543^{+0.024}_{-0.022}$	35^{+4}_{-4}	$1.61^{+0.12}_{-0.10}$	$0.25^{+0.02}_{-0.02}$	49^{+12}_{-11}
A3667	0.0560	$7.00^{+0.60}_{-0.60}$	$0.541^{+0.008}_{-0.008}$	279^{+10}_{-10}	$0.33^{+0.01}_{-0.01}$	$2.81^{+0.21}_{-0.17}$	0^{+0}_{-0}
A4038	0.0283	$3.15^{+0.03}_{-0.03}$	$0.541^{+0.008}_{-0.008}$	58^{+3}_{-3}	$1.49^{+0.09}_{-0.09}$	$0.40^{+0.02}_{-0.02}$	68^{+14}_{-12}
A4059	0.0460	$4.10^{+0.30}_{-0.30}$	$0.582^{+0.010}_{-0.010}$	89^{+5}_{-5}	$1.18^{+0.08}_{-0.08}$	$0.58^{+0.05}_{-0.05}$	69^{+20}_{-15}
COMA	0.0232	$8.38^{+0.34}_{-0.34}$	$0.654^{+0.019}_{-0.021}$	343^{+22}_{-20}	$0.30^{+0.06}_{-0.06}$	$3.45^{+0.85}_{-0.56}$	0^{+0}_{-0}
EXO0422	0.0390	$2.90^{+0.90}_{-0.60}$	$0.722^{+0.104}_{-0.071}$	142^{+40}_{-30}	$0.66^{+0.11}_{-0.08}$	$0.85^{+0.23}_{-0.19}$	48^{+59}_{-38}
FORNAX	0.0046	$1.20^{+0.04}_{-0.04}$	$0.804^{+0.098}_{-0.084}$	173^{+17}_{-15}	$0.09^{+0.01}_{-0.01}$	$2.50^{+0.20}_{-0.18}$	0^{+0}_{-0}
HYDRA-A	0.0538	$3.80^{+0.20}_{-0.20}$	$0.573^{+0.003}_{-0.003}$	50^{+1}_{-1}	$3.58^{+0.37}_{-0.37}$	$0.18^{+0.02}_{-0.02}$	293^{+150}_{-84}
IIIZw54	0.0311	$(2.16^{+0.35}_{-0.30})$	$0.887^{+0.320}_{-0.151}$	289^{+123}_{-72}	$0.20^{+0.03}_{-0.03}$	$2.31^{+0.55}_{-0.56}$	0^{+0}_{-0}
MKW3S	0.0450	$3.50^{+0.20}_{-0.20}$	$0.581^{+0.008}_{-0.007}$	66^{+2}_{-2}	$1.89^{+0.25}_{-0.25}$	$0.33^{+0.06}_{-0.04}$	121^{+76}_{-44}
MKW4	0.0200	$1.71^{+0.09}_{-0.09}$	$0.440^{+0.004}_{-0.005}$	11^{+0}_{-0}	$2.92^{+0.11}_{-0.09}$	$0.13^{+0.01}_{-0.01}$	16^{+2}_{-2}
MKW8	0.0270	$3.29^{+0.23}_{-0.22}$	$0.511^{+0.098}_{-0.059}$	106^{+70}_{-42}	$0.26^{+0.11}_{-0.05}$	$2.31^{+0.62}_{-0.69}$	0^{+0}_{-0}
NGC1550	0.0123	$1.43^{+0.04}_{-0.03}$	$0.554^{+0.049}_{-0.037}$	45^{+15}_{-10}	$0.75^{+0.14}_{-0.09}$	$0.41^{+0.06}_{-0.06}$	20^{+10}_{-8}
NGC4636	0.0037	$0.76^{+0.01}_{-0.01}$	$0.491^{+0.032}_{-0.027}$	6^{+2}_{-1}	$1.68^{+0.49}_{-0.24}$	$0.07^{+0.01}_{-0.01}$	2^{+2}_{-1}
NGC5044	0.0090	$1.07^{+0.01}_{-0.01}$	$0.524^{+0.002}_{-0.003}$	11^{+0}_{-0}	$3.45^{+0.03}_{-0.03}$	$0.05^{+0.00}_{-0.00}$	28^{+1}_{-1}
NGC507	0.0165	$1.26^{+0.07}_{-0.07}$	$0.444^{+0.005}_{-0.005}$	19^{+1}_{-1}	$1.16^{+0.04}_{-0.04}$	$0.22^{+0.01}_{-0.01}$	14^{+2}_{-2}
S1101	0.0580	$3.00^{+1.20}_{-0.70}$	$0.639^{+0.006}_{-0.007}$	55^{+1}_{-1}	$2.90^{+0.19}_{-0.18}$	$0.20^{+0.06}_{-0.04}$	299^{+179}_{-112}
ZwCl1215	0.0750	$(5.58^{+0.89}_{-0.78})$	$0.819^{+0.038}_{-0.034}$	431^{+27}_{-25}	$0.27^{+0.01}_{-0.01}$	$3.00^{+0.36}_{-0.33}$	0^{+0}_{-0}

Note: ^{a)}: Markevitch 1998. ^{b)}: White 2000. ^{c)}: Edge & Stewart 1991. T_m with a, b or c has no cooling flow correction. ^{d)}: Fukazawa et al. 1998 with cooling flow correction. T_m in a bracket is estimated from the L_X - T relation given by Markevitch (1998). Others are from Reiprich & Böhringer (2002) and references therein.

Table 2. Continued (extended cluster sample).

Name	z	T_m (keV)	β	r_c (kpc)	n_{center} (10^{-2} cm $^{-3}$)	t_{cool} (10^{10} yr)	\dot{M} (M_{\odot}/yr)
3C129	0.0223	$^c 5.60^{+0.70}_{-0.60}$	$0.601^{+0.260}_{-0.131}$	318^{+178}_{-107}	$0.18^{+0.07}_{-0.04}$	$4.71^{+1.62}_{-1.39}$	0^{+0}_{-0}
A0539	0.0288	$^d 3.24^{+0.09}_{-0.09}$	$0.561^{+0.020}_{-0.018}$	147^{+13}_{-12}	$0.72^{+0.12}_{-0.09}$	$0.84^{+0.13}_{-0.12}$	3^{+1}_{-1}
A0548e	0.0410	$^b 3.10^{+0.10}_{-0.10}$	$0.480^{+0.013}_{-0.013}$	118^{+12}_{-11}	$0.28^{+0.02}_{-0.01}$	$2.09^{+0.11}_{-0.12}$	0^{+0}_{-0}
A0548w	0.0424	$(1.20^{+0.19}_{-0.17})$	$0.666^{+0.194}_{-0.111}$	198^{+89}_{-61}	$0.10^{+0.03}_{-0.02}$	$2.28^{+0.67}_{-0.59}$	0^{+0}_{-29}
A0644	0.0704	$^a 7.10^{+0.60}_{-0.26}$	$0.700^{+0.011}_{-0.017}$	202^{+6}_{-6}	$0.78^{+0.01}_{-0.01}$	$1.18^{+0.08}_{-0.08}$	16^{+15}_{-15}
A1413	0.1427	$^b 7.32^{+0.26}_{-0.24}$	$0.660^{+0.011}_{-0.015}$	178^{+12}_{-11}	$1.24^{+0.07}_{-0.07}$	$0.72^{+0.05}_{-0.04}$	190^{+40}_{-32}
A1689	0.1840	$^b 9.23^{+0.28}_{-0.28}$	$0.690^{+0.011}_{-0.011}$	162^{+6}_{-6}	$2.12^{+0.18}_{-0.18}$	$0.47^{+0.04}_{-0.04}$	683^{+239}_{-182}
A1775	0.0757	$^b 3.69^{+0.20}_{-0.11}$	$0.673^{+0.026}_{-0.023}$	259^{+19}_{-17}	$0.30^{+0.05}_{-0.04}$	$2.16^{+0.36}_{-0.28}$	0^{+0}_{-0}
A1800	0.0748	$(4.02^{+0.64}_{-0.56})$	$0.766^{+0.308}_{-0.139}$	391^{+223}_{-131}	$0.18^{+0.07}_{-0.04}$	$3.65^{+1.14}_{-1.07}$	0^{+0}_{-0}
A1914	0.1712	$^b 10.53^{+0.51}_{-0.50}$	$0.751^{+0.018}_{-0.017}$	230^{+10}_{-10}	$1.12^{+0.03}_{-0.03}$	$0.97^{+0.04}_{-0.04}$	180^{+50}_{-40}
A2151w	0.0369	$^b 2.40^{+0.06}_{-0.06}$	$0.564^{+0.014}_{-0.013}$	68^{+5}_{-4}	$0.82^{+0.04}_{-0.03}$	$0.60^{+0.03}_{-0.03}$	30^{+3}_{-3}
A2319	0.0564	$^a 9.20^{+0.70}_{-0.70}$	$0.591^{+0.012}_{-0.012}$	284^{+14}_{-13}	$0.51^{+0.05}_{-0.05}$	$2.10^{+0.24}_{-0.21}$	0^{+0}_{-0}
A2734	0.0620	$(3.85^{+0.62}_{-0.54})$	$0.624^{+0.034}_{-0.029}$	211^{+26}_{-23}	$0.32^{+0.02}_{-0.02}$	$2.04^{+0.28}_{-0.27}$	0^{+0}_{-0}
A2877	0.0241	$3.50^{+2.20}_{-1.10}$	$0.566^{+0.029}_{-0.025}$	189^{+18}_{-16}	$0.19^{+0.02}_{-0.02}$	$3.33^{+1.60}_{-0.95}$	0^{+0}_{-0}
A3395n	0.0498	$^a 4.80^{+0.40}_{-0.40}$	$0.981^{+0.619}_{-0.244}$	672^{+383}_{-203}	$0.10^{+0.04}_{-0.02}$	$7.33^{+2.40}_{-2.03}$	0^{+0}_{-0}
A3528n	0.0540	$3.40^{+1.66}_{-0.64}$	$0.621^{+0.034}_{-0.030}$	177^{+16}_{-15}	$0.34^{+0.02}_{-0.02}$	$1.81^{+0.61}_{-0.28}$	0^{+0}_{-0}
A3528s	0.0551	$3.15^{+0.89}_{-0.59}$	$0.463^{+0.013}_{-0.012}$	100^{+8}_{-8}	$0.48^{+0.03}_{-0.02}$	$1.21^{+0.26}_{-0.19}$	1^{+1}_{-1}
A3530	0.0544	$3.89^{+0.27}_{-0.25}$	$0.773^{+0.114}_{-0.085}$	420^{+74}_{-61}	$0.13^{+0.01}_{-0.01}$	$5.27^{+0.61}_{-0.56}$	0^{+0}_{-0}
A3532	0.0539	$4.58^{+0.19}_{-0.17}$	$0.653^{+0.034}_{-0.029}$	281^{+26}_{-24}	$0.30^{+0.06}_{-0.05}$	$2.42^{+0.50}_{-0.38}$	0^{+0}_{-0}
A3560	0.0495	$(3.16^{+0.51}_{-0.44})$	$0.566^{+0.033}_{-0.029}$	255^{+30}_{-27}	$0.17^{+0.01}_{-0.01}$	$3.47^{+0.51}_{-0.46}$	0^{+0}_{-0}
A3627	0.0163	$^b 6.02^{+0.08}_{-0.08}$	$0.555^{+0.056}_{-0.044}$	299^{+56}_{-49}	$0.19^{+0.02}_{-0.02}$	$4.51^{+0.44}_{-0.45}$	0^{+0}_{-0}
A3695	0.0890	$(5.29^{+0.85}_{-0.74})$	$0.642^{+0.259}_{-0.117}$	398^{+253}_{-149}	$0.20^{+0.08}_{-0.05}$	$3.82^{+1.27}_{-1.12}$	0^{+0}_{-0}
A3822	0.0760	$(4.90^{+0.78}_{-0.78})$	$0.639^{+0.098}_{-0.086}$	350^{+159}_{-111}	$0.21^{+0.05}_{-0.05}$	$3.60^{+0.83}_{-0.86}$	0^{+0}_{-0}
A3827	0.0980	$(7.08^{+1.13}_{-0.99})$	$0.989^{+0.110}_{-0.192}$	593^{+247}_{-148}	$0.26^{+0.06}_{-0.05}$	$3.43^{+0.86}_{-0.72}$	0^{+0}_{-0}
A3888	0.1510	$(8.84^{+1.41}_{-1.24})$	$0.928^{+0.084}_{-0.066}$	400^{+45}_{-39}	$0.52^{+0.04}_{-0.03}$	$1.92^{+0.26}_{-0.25}$	0^{+0}_{-0}
A3921	0.0936	$^b 5.73^{+0.24}_{-0.23}$	$0.762^{+0.036}_{-0.030}$	328^{+25}_{-23}	$0.34^{+0.01}_{-0.01}$	$2.34^{+0.12}_{-0.12}$	0^{+0}_{-0}
AWM7	0.0172	$^d 3.73^{+0.09}_{-0.09}$	$0.671^{+0.027}_{-0.025}$	173^{+17}_{-15}	$0.60^{+0.05}_{-0.05}$	$1.10^{+0.11}_{-0.09}$	6^{+3}_{-4}
HCG94	0.0417	$3.45^{+0.30}_{-0.30}$	$0.514^{+0.007}_{-0.006}$	86^{+4}_{-3}	$0.70^{+0.02}_{-0.02}$	$0.89^{+0.07}_{-0.07}$	6^{+3}_{-2}
II Zw 108	0.0494	$(3.44^{+0.53}_{-0.53})$	$0.662^{+0.167}_{-0.048}$	365^{+159}_{-105}	$0.14^{+0.04}_{-0.02}$	$4.50^{+1.06}_{-1.04}$	0^{+0}_{-0}
M49	0.0044	$0.95^{+0.02}_{-0.01}$	$0.592^{+0.007}_{-0.007}$	10^{+0}_{-0}	$1.33^{+0.02}_{-0.02}$	$0.12^{+0.00}_{-0.00}$	2^{+0}_{-0}
NGC 499	0.0147	$0.72^{+0.03}_{-0.02}$	$0.722^{+0.034}_{-0.030}$	23^{+2}_{-1}	$0.94^{+0.10}_{-0.09}$	$0.12^{+0.01}_{-0.01}$	11^{+4}_{-3}
NGC 5813	0.0064	$(0.52^{+0.08}_{-0.07})$	$0.766^{+0.179}_{-0.103}$	24^{+8}_{-5}	$0.90^{+0.18}_{-0.13}$	$0.10^{+0.02}_{-0.02}$	9^{+11}_{-5}
NGC 5846	0.0061	$0.82^{+0.01}_{-0.01}$	$0.599^{+0.016}_{-0.015}$	6^{+0}_{-0}	$3.56^{+0.22}_{-0.21}$	$0.04^{+0.00}_{-0.00}$	2^{+0}_{-0}
OPHIUCHU	0.0280	$10.26^{+0.32}_{-0.32}$	$0.747^{+0.035}_{-0.032}$	278^{+23}_{-21}	$0.68^{+0.04}_{-0.04}$	$1.72^{+0.12}_{-0.10}$	0^{+0}_{-0}
PERSEUS	0.0183	$^d 6.79^{+0.12}_{-0.12}$	$0.540^{+0.006}_{-0.006}$	63^{+2}_{-2}	$3.25^{+0.06}_{-0.06}$	$0.28^{+0.01}_{-0.01}$	481^{+31}_{-32}
PKS 0745	0.1028	$^b 7.21^{+0.11}_{-0.11}$	$0.608^{+0.006}_{-0.006}$	71^{+2}_{-2}	$5.70^{+0.04}_{-0.15}$	$0.16^{+0.00}_{-0.00}$	1424^{+150}_{-133}
RXJ2344	0.0786	$(4.73^{+0.76}_{-0.66})$	$0.807^{+0.033}_{-0.030}$	300^{+19}_{-18}	$0.51^{+0.06}_{-0.05}$	$1.43^{+0.24}_{-0.20}$	0^{+0}_{-0}
S405	0.0613	$(4.21^{+0.67}_{-0.59})$	$0.664^{+0.263}_{-0.133}$	458^{+261}_{-158}	$0.12^{+0.05}_{-0.03}$	$5.67^{+1.85}_{-1.74}$	0^{+0}_{-0}
S540	0.0358	$(2.40^{+0.38}_{-0.34})$	$0.641^{+0.073}_{-0.051}$	130^{+38}_{-28}	$0.40^{+0.07}_{-0.05}$	$1.23^{+0.23}_{-0.24}$	1^{+7}_{-1}
S636	0.0116	$(1.18^{+0.19}_{-0.17})$	$0.752^{+0.217}_{-0.123}$	343^{+130}_{-86}	$0.07^{+0.01}_{-0.01}$	$3.06^{+0.77}_{-0.65}$	0^{+0}_{-0}
TRIANGUL	0.0510	$^a 9.50^{+0.70}_{-0.70}$	$0.610^{+0.010}_{-0.010}$	278^{+10}_{-9}	$0.53^{+0.01}_{-0.01}$	$1.98^{+0.12}_{-0.12}$	0^{+0}_{-0}
UGC 03957	0.0340	$(2.58^{+0.41}_{-0.36})$	$0.740^{+0.133}_{-0.086}$	142^{+44}_{-33}	$0.48^{+0.08}_{-0.06}$	$1.09^{+0.21}_{-0.20}$	8^{+11}_{-8}
ZwCl1742	0.0757	$(5.23^{+0.84}_{-0.73})$	$0.717^{+0.073}_{-0.053}$	231^{+45}_{-38}	$0.60^{+0.11}_{-0.09}$	$1.29^{+0.28}_{-0.23}$	0^{+23}_{-0}

Note: The symbols in T_m have the same meanings as that in Table 1.

such that the overall effect may remain undetected (see also discussion below).

Among the relations listed in Table 7, the $M_{500}-T_h$ relation, together with the $M_{gas,500}-T_h$ relation (Fig. 11), has the smallest scatter. This could in part be due to the fact that the mass and temperature parameter are correlated. We will show below (Sect. 4.5) by means of the $M_{gas,500}-T_h$ relation that this is not a large effect and thus not the main reason for the different L_X normalization. Rather the tight relation shows that mass and temperature are linked in a more fundamental way by simple self-similar gravitational processes than the other relations.

4.3. L_X-T relation

The L_X-T_m and L_X-T_h relations for the cluster sample with measured T_m and T_h derived from ASCA are shown in Figs. 6 and 7. The values of L_X are the X-ray luminosities in (0.1-2.4 keV) derived from Ikebe et al. (2002). Since for some clusters T_m is measured with the central region excluded (cooling flow correction), here we only use T_m without cooling flow correction. For T_m derived from Fukazawa et al. (1998) which includes cooling flow correction, we use the central T_m ($0-2' \sim 3'$) instead (Fukazawa et al. 2000). The resultant slopes of L_X-T relations are, 2.23 ± 0.15 and 2.73 ± 0.13 for T_m and T_h , respectively.

Table 3. Fit parameters with a double β model. χ_s^2 and χ_d^2 are the reduced χ^2 values for the single β model and the double β model, respectively. S_1 and S_2 are the surface brightness. Note $1' = 120$ pixels.

Name	S_{01} (10^{-6} cts/s/pixel 2)	S_{02} (10^{-6} cts/s/pixel 2)	β_1	β_2	r_{c1} (kpc)	r_{c2} (kpc)	χ_s^2	χ_d^2
A0085	27.01±0.99	1.70±0.24	0.60±0.03	0.73±0.03	58±3	385±28	5.32	1.90
A0119	0.50±0.03	0.16±0.03	0.76±0.11	1.46±0.24	399±45	1511±207	1.75	1.47
A0133	23.96±0.82	0.87±0.06	0.65±0.02	0.78±0.02	42±1	321±13	4.14	1.71
A0401	3.93±0.30	0.51±0.30	0.69±0.08	0.66±0.03	239±23	525±93	1.19	1.17
A0478	37.46±1.30	4.02±0.71	0.68±0.04	0.71±0.02	72±4	253±14	3.53	1.50
A0496	37.20±1.87	1.40±0.33	0.59±0.05	0.69±0.04	30±3	257±31	3.97	1.07
A1367	0.30±0.01	0.22±0.01	0.96±0.10	1.51±0.06	291±23	976±31	1.51	1.14
A1644	1.16±0.24	0.18±0.06	0.83±0.30	2.38±0.81	274±92	2169±532	0.97	0.89
A1651	7.39±0.47	2.17±0.24	0.75±0.07	0.76±0.02	120±11	356±17	1.46	1.10
A1795	41.44±0.60	1.75±0.16	0.72±0.02	0.89±0.02	79±2	432±19	10.04	1.99
A2029	55.67±1.98	5.81±1.16	0.63±0.03	0.65±0.02	62±3	213±15	2.61	1.40
A2052	23.26±1.43	3.78±0.46	2.10±1.02	0.66±0.02	84±25	140±12	2.70	1.45
A2063	4.71±0.00	0.51±0.00	0.49±0.00	2.02±0.00	55±0	640±0	1.97	1.17
A2142	19.59±0.69	0.86±0.19	0.67±0.04	1.01±0.13	140±8	893±130	2.55	1.39
A2255	0.35±0.05	0.56±0.05	1.15±0.32	0.90±0.03	469±93	778±31	1.38	1.36
A2589	3.45±0.28	0.51±0.23	0.66±0.10	0.74±0.07	95±13	311±56	1.38	1.23
A2634	0.42±0.04	0.13±0.01	0.47±0.03	1.89±0.18	80±10	1189±77	1.77	1.25
A2657	2.34±0.00	0.25±0.00	0.89±0.00	1.27±0.00	148±0	796±0	3.05	1.68
A3112	41.34±2.59	3.11±1.41	0.63±0.06	0.62±0.03	51±5	164±24	1.62	1.44
A3266	1.24±0.05	0.76±0.04	1.20±0.11	1.27±0.03	450±29	1162±28	2.62	1.76
A3391	0.87±0.16	0.66±0.08	0.50±0.06	0.66±0.03	71±19	335±24	1.30	1.26
A3526	8.97±0.79	0.34±0.07	0.57±0.03	0.70±0.05	33±3	272±32	1.22	1.11
A3558	3.13±0.05	0.20±0.04	0.68±0.03	1.17±0.11	232±8	1198±119	3.66	2.90
A3562	2.84±0.10	0.07±0.03	0.52±0.02	1.26±0.27	103±6	1341±273	2.10	1.78
A3571	5.00±0.40	4.39±0.22	0.82±0.13	0.68±0.01	94±13	256±6	1.72	1.13
A3667	2.10±0.04	0.20±0.01	0.89±0.03	1.70±0.09	402±12	2375±95	2.53	1.63
A4038	10.37±0.58	0.49±0.29	0.58±0.04	0.70±0.12	53±4	241±69	1.27	1.18
A4059	8.67±0.46	0.43±0.18	0.64±0.06	0.90±0.19	82±7	438±112	1.32	1.06
COMA	1.13±0.54	1.35±0.52	0.57±0.05	0.90±0.23	329±66	444±104	1.09	1.09
HYDRA-A	40.63±1.40	5.67±0.43	1.84±0.42	0.73±0.01	98±14	183±8	6.24	1.81
MKW3S	12.40±0.92	3.84±0.62	1.42±0.41	0.68±0.02	91±17	152±13	2.07	1.54
S1101	34.57±1.35	0.58±0.30	0.79±0.08	0.96±0.16	66±6	381±104	1.67	1.33
A0539	1.53±0.19	0.41±0.08	0.53±0.09	0.75±0.08	42±9	313±40	1.77	1.19
A1413	10.29±0.59	1.11±0.20	0.80±0.07	0.91±0.05	155±12	559±47	1.73	1.45
A1689	23.46±1.39	2.53±0.82	0.88±0.14	0.91±0.05	152±19	471±56	1.73	1.24
A1775	1.48±0.08	0.18±0.04	2.05±0.69	1.70±0.53	540±114	1443±382	1.62	1.46
A2319	3.59±0.23	0.76±0.19	1.06±0.19	0.82±0.06	383±48	874±120	1.34	1.24
A2877	0.28±0.02	0.15±0.01	3.58±0.89	1.23±0.07	323±47	606±28	2.79	2.04
A3532	0.91±0.11	0.27±0.09	0.74±0.24	1.09±0.21	193±48	761±161	1.31	1.12
A3888	3.90±0.22	0.30±0.15	1.39±0.19	1.71±0.39	503±47	1341±271	0.90	0.90
AWM7	2.53±0.22	0.56±0.11	0.78±0.11	0.88±0.05	125±16	406±39	1.40	1.33
HCG94	2.26±0.09	0.41±0.04	0.53±0.01	0.58±0.01	59±3	199±10	2.43	2.20
NGC507	1.20±0.30	3.37±0.34	0.76±0.04	4.29±1.21	41±4	73±11	2.41	2.40
NGC5846	10.88±0.97	1.48±0.18	0.51±0.01	4.78±1.07	3±0	55±7	1.78	1.53
OPHIUCHU	4.86±0.24	0.35±0.10	1.04±0.11	1.40±0.22	328±26	1190±177	1.41	1.37
PKS0745	53.06±1.73	2.02±0.39	0.70±0.02	0.65±0.01	72±2	235±16	1.17	1.06
RXJ2344	1.41±0.20	1.41±0.14	0.72±0.13	0.92±0.04	128±27	400±20	1.76	1.70
TRIANGUL	3.36±0.09	0.66±0.06	0.71±0.03	0.80±0.02	248±9	700±30	1.57	1.28
ZwCl1742	3.50±0.57	0.58±0.33	0.83±0.21	0.93±0.16	194±45	547±135	1.11	1.11

They are much higher than 1.5, predicted by self-similar model for the L_X - T relation (note L_X is in the ROSAT band, not bolometric). This is consistent with the results in Reiprich & Böhringer (2002). Note that the slope of L_X - T_m is shallower than that of L_X - T_h . This may be due to the facts that T_m has an offset to the low temperature direction compared to T_h and fewer low L_X clusters are included in the T_m relation. Remarkable is the clearly higher normalization of the relation for CCCs compared to NCCCs, with offsets of factors of 2.05 and 1.84 for the relation with T_m and T_h , respectively. The normalization difference is slightly small using T_h in the scaling relation and the scatter is also slightly reduced, since most probably T_h provides a better measure of the global gravitational potential depth and better mass proxy, as this temperature is not so much downward biased by the central cooling core region as T_m . But the difference between the two relations is not very large. This is already an in-

dication that biased temperature measurements for CCCs are not the major reason for the different normalizations of the relations for CCCs and NCCCs. Here the effect of the different mass coverage of CCCs and NCCCs on the slope of the combined relation (which should make the relation shallower) is not so strong as in Fig. 4. The combined relation is slightly shallower than the NCCC and CCC relations.

4.4. L_X - M relation

The L_X - M relation is the most important relation for the application to cosmological cluster surveys. The previously determined relation for this sample (Reiprich & Böhringer, 2002) was used to get cluster mass estimates for example for the cosmological studies (Schuecker et al. 2001b, 2003 and Stanek et al. 2006). The resultant slope of

Table 4. Cluster properties for the relations.

Name	T_h (keV)	L_X (10^{44} erg/s, 0.1-2.4keV)	M_{500} ($10^{14} M_\odot$)	r_{500} (Mpc)	f_{gas}	$M_{gas,500}$ ($10^{13} M_\odot$)	L_{nir} ($h_{70}^{-2} 10^{12} L_\odot$)
2A0335	$3.64^{+0.09}_{-0.08}$	4.64 ± 0.04	$2.79^{+1.09}_{-1.63}$	$1.20^{+0.14}_{-0.30}$	$0.19^{+0.13}_{-0.03}$	$5.42^{+0.80}_{-1.74}$	3.44 ± 0.07
A0085	$6.51^{+0.16}_{-0.23}$	9.67 ± 0.11	$8.08^{+1.37}_{-3.51}$	$1.68^{+0.10}_{-0.29}$	$0.21^{+0.09}_{-0.02}$	$16.66^{+1.49}_{-3.54}$	5.99 ± 0.06
A0119	$5.69^{+0.24}_{-0.28}$	3.34 ± 0.05	$8.98^{+1.20}_{-2.59}$	$1.76^{+0.07}_{-0.19}$	$0.15^{+0.04}_{-0.01}$	$13.24^{+0.75}_{-1.79}$	6.46 ± 0.07
A0133	$3.97^{+0.28}_{-0.27}$	2.85 ± 0.04	$4.30^{+1.00}_{-2.08}$	$1.36^{+0.10}_{-0.27}$	$0.14^{+0.07}_{-0.02}$	$6.01^{+0.51}_{-1.36}$	3.72 ± 0.04
A0262	$2.25^{+0.66}_{-0.96}$	0.98 ± 0.06	$0.94^{+0.05}_{-0.13}$	$0.85^{+0.02}_{-0.04}$	$0.21^{+0.05}_{-0.05}$	$1.93^{+0.39}_{-0.48}$	2.07 ± 0.07
A0399	$6.46^{+0.38}_{-0.36}$	7.13 ± 0.62	$7.74^{+0.15}_{-2.32}$	$1.63^{+0.04}_{-0.18}$	$0.19^{+0.05}_{-0.07}$	$14.79^{+5.43}_{-5.62}$	-
A0400	$2.43^{+0.13}_{-0.12}$	0.65 ± 0.01	$1.33^{+0.11}_{-0.16}$	$0.95^{+0.02}_{-0.04}$	$0.16^{+0.02}_{-0.01}$	$2.11^{+0.13}_{-0.20}$	2.34 ± 0.06
A0401	$7.19^{+0.28}_{-0.24}$	12.41 ± 0.22	$8.38^{+1.22}_{-2.84}$	$1.67^{+0.08}_{-0.21}$	$0.25^{+0.08}_{-0.02}$	$20.55^{+2.75}_{-4.27}$	-
A0478	$6.91^{+0.40}_{-0.36}$	17.44 ± 0.18	$8.85^{+3.00}_{-4.69}$	$1.68^{+0.17}_{-0.37}$	$0.22^{+0.14}_{-0.04}$	$19.10^{+2.18}_{-4.72}$	7.96 ± 0.09
A0496	$4.59^{+0.10}_{-0.10}$	3.77 ± 0.05	$4.81^{+0.89}_{-2.11}$	$1.44^{+0.08}_{-0.25}$	$0.16^{+0.08}_{-0.02}$	$7.77^{+0.77}_{-1.65}$	3.91 ± 0.05
A0576	$3.83^{+0.16}_{-0.15}$	1.86 ± 0.21	$4.61^{+3.25}_{-2.39}$	$1.42^{+0.28}_{-0.30}$	$0.12^{+0.18}_{-0.08}$	$5.43^{+2.11}_{-3.53}$	4.62 ± 0.06
A0754	$9.00^{+0.34}_{-0.34}$	3.97 ± 0.11	$13.86^{+4.40}_{-6.43}$	$2.02^{+0.19}_{-0.38}$	$0.10^{+0.05}_{-0.02}$	$13.25^{+1.83}_{-3.32}$	9.28 ± 0.05
A1060	$3.15^{+0.05}_{-0.05}$	0.56 ± 0.03	$2.50^{+0.62}_{-1.02}$	$1.19^{+0.09}_{-0.19}$	$0.09^{+0.04}_{-0.02}$	$2.19^{+0.49}_{-0.69}$	2.39 ± 0.06
A1367	$3.55^{+0.08}_{-0.08}$	1.20 ± 0.02	$7.42^{+1.11}_{-2.37}$	$1.69^{+0.08}_{-0.20}$	$0.09^{+0.03}_{-0.01}$	$6.39^{+0.30}_{-0.69}$	3.81 ± 0.05
A1644	$4.70^{+0.90}_{-0.70}$	3.92 ± 0.34	$7.34^{+4.30}_{-4.40}$	$1.64^{+0.27}_{-0.43}$	$0.17^{+0.13}_{-0.04}$	$12.53^{+2.71}_{-4.28}$	5.82 ± 0.05
A1650	$5.68^{+0.30}_{-0.27}$	7.33 ± 0.79	$6.53^{+2.17}_{-2.43}$	$1.52^{+0.15}_{-0.22}$	$0.19^{+0.14}_{-0.08}$	$12.22^{+4.85}_{-5.79}$	3.09 ± 0.14
A1651	$6.22^{+0.41}_{-0.22}$	7.85 ± 0.14	$8.29^{+1.95}_{-2.90}$	$1.65^{+0.12}_{-0.28}$	$0.17^{+0.08}_{-0.02}$	$14.38^{+2.81}_{-2.81}$	7.82 ± 0.05
A1736	$3.68^{+0.22}_{-0.17}$	3.22 ± 0.33	$2.17^{+0.62}_{-0.69}$	$1.09^{+0.10}_{-0.13}$	$0.25^{+0.09}_{-0.10}$	$5.33^{+1.07}_{-1.84}$	5.28 ± 0.10
A1795	$6.17^{+0.26}_{-0.25}$	10.00 ± 0.07	$9.87^{+3.85}_{-5.48}$	$1.79^{+0.21}_{-0.42}$	$0.15^{+0.11}_{-0.03}$	$14.51^{+1.50}_{-3.46}$	4.80 ± 0.04
A2029	$7.93^{+0.39}_{-0.36}$	17.07 ± 0.18	$9.95^{+3.29}_{-5.16}$	$1.77^{+0.18}_{-0.38}$	$0.21^{+0.12}_{-0.03}$	$20.92^{+2.45}_{-5.52}$	8.12 ± 0.06
A2052	$3.12^{+0.10}_{-0.09}$	2.37 ± 0.04	$2.70^{+0.57}_{-1.32}$	$1.19^{+0.08}_{-0.24}$	$0.16^{+0.09}_{-0.02}$	$4.31^{+1.43}_{-0.95}$	3.22 ± 0.05
A2063	$3.56^{+0.16}_{-0.12}$	2.26 ± 0.05	$2.36^{+0.24}_{-0.59}$	$1.14^{+0.04}_{-0.10}$	$0.19^{+0.03}_{-0.01}$	$4.44^{+0.20}_{-0.52}$	3.50 ± 0.04
A2065	$5.37^{+0.34}_{-0.30}$	5.63 ± 0.55	$11.19^{+9.87}_{-8.82}$	$1.84^{+0.42}_{-0.50}$	$0.12^{+0.08}_{-0.01}$	$13.68^{+5.07}_{-9.46}$	7.33 ± 0.05
A2142	$8.46^{+0.53}_{-0.49}$	21.05 ± 0.29	$14.33^{+3.64}_{-6.83}$	$1.97^{+0.15}_{-0.38}$	$0.22^{+0.11}_{-0.03}$	$31.94^{+3.25}_{-7.17}$	7.20 ± 0.09
A2147	$4.34^{+0.12}_{-0.13}$	2.87 ± 0.15	$2.31^{+0.40}_{-0.36}$	$1.13^{+0.06}_{-0.06}$	$0.26^{+0.07}_{-0.08}$	$5.93^{+1.19}_{-1.66}$	4.15 ± 0.05
A2163	$10.55^{+1.01}_{-0.68}$	32.16 ± 0.82	$16.00^{+3.48}_{-4.86}$	$1.85^{+0.13}_{-0.21}$	$0.27^{+0.08}_{-0.03}$	$43.75^{+3.73}_{-7.16}$	-
A2199	$4.28^{+0.10}_{-0.10}$	4.20 ± 0.12	$4.29^{+1.18}_{-1.89}$	$1.39^{+0.12}_{-0.24}$	$0.17^{+0.08}_{-0.03}$	$7.35^{+0.91}_{-1.73}$	4.43 ± 0.04
A2204	$6.38^{+0.23}_{-0.23}$	25.89 ± 0.69	$5.82^{+1.91}_{-2.98}$	$1.38^{+0.14}_{-0.29}$	$0.29^{+0.15}_{-0.05}$	$16.85^{+2.24}_{-4.70}$	-
A2244	$5.77^{+0.61}_{-0.44}$	8.30 ± 0.28	$5.48^{+2.48}_{-2.23}$	$1.42^{+0.22}_{-0.23}$	$0.21^{+0.08}_{-0.04}$	$11.52^{+1.71}_{-2.65}$	-
A2255	$5.92^{+0.40}_{-0.26}$	5.46 ± 0.11	$7.86^{+0.92}_{-1.67}$	$1.63^{+0.06}_{-0.12}$	$0.19^{+0.03}_{-0.01}$	$14.66^{+1.06}_{-1.50}$	8.70 ± 0.08
A2256	$6.83^{+0.23}_{-0.21}$	9.24 ± 0.22	$12.12^{+3.41}_{-4.12}$	$1.91^{+0.16}_{-0.25}$	$0.18^{+0.07}_{-0.03}$	$21.78^{+2.19}_{-4.09}$	10.11 ± 0.04
A2589	$3.38^{+0.13}_{-0.13}$	1.87 ± 0.04	$3.24^{+0.54}_{-1.40}$	$1.26^{+0.07}_{-0.21}$	$0.15^{+0.08}_{-0.02}$	$4.76^{+0.81}_{-0.99}$	2.51 ± 0.06
A2597	$4.20^{+0.49}_{-0.41}$	6.75 ± 0.14	$3.71^{+1.35}_{-2.22}$	$1.26^{+0.14}_{-0.33}$	$0.17^{+0.13}_{-0.03}$	$6.48^{+0.86}_{-2.02}$	-
A2634	$3.45^{+0.16}_{-0.16}$	0.99 ± 0.03	$4.51^{+0.67}_{-1.00}$	$1.42^{+0.07}_{-0.11}$	$0.11^{+0.02}_{-0.01}$	$5.10^{+0.36}_{-0.37}$	4.53 ± 0.08
A2657	$3.53^{+0.12}_{-0.12}$	1.74 ± 0.03	$6.06^{+0.92}_{-2.57}$	$1.55^{+0.11}_{-0.26}$	$0.10^{+0.05}_{-0.01}$	$6.26^{+0.35}_{-1.08}$	2.35 ± 0.09
A3112	$4.72^{+0.37}_{-0.25}$	7.32 ± 0.15	$4.36^{+1.26}_{-2.25}$	$1.34^{+0.12}_{-0.29}$	$0.20^{+0.13}_{-0.03}$	$8.87^{+1.71}_{-2.49}$	4.11 ± 0.08
A3158	$5.41^{+0.26}_{-0.24}$	5.61 ± 0.15	$5.75^{+0.89}_{-1.66}$	$1.49^{+0.07}_{-0.16}$	$0.20^{+0.05}_{-0.03}$	$11.63^{+1.13}_{-2.10}$	6.76 ± 0.05
A3266	$7.72^{+0.35}_{-0.28}$	8.62 ± 0.11	$19.24^{+4.76}_{-7.58}$	$2.23^{+0.17}_{-0.34}$	$0.14^{+0.05}_{-0.02}$	$27.35^{+1.97}_{-4.39}$	7.79 ± 0.06
A3376	$4.43^{+0.39}_{-0.38}$	2.16 ± 0.05	$6.77^{+1.55}_{-1.99}$	$1.60^{+0.11}_{-0.17}$	$0.13^{+0.04}_{-0.02}$	$8.70^{+0.86}_{-1.50}$	3.81 ± 0.04
A3391	$5.89^{+0.45}_{-0.39}$	2.64 ± 0.08	$6.04^{+0.74}_{-1.69}$	$1.53^{+0.10}_{-0.16}$	$0.14^{+0.05}_{-0.01}$	$8.63^{+0.52}_{-1.01}$	5.84 ± 0.09
A3395s	$5.55^{+0.89}_{-0.65}$	2.12 ± 0.13	$9.48^{+4.35}_{-4.29}$	$1.78^{+0.24}_{-0.32}$	$0.10^{+0.09}_{-0.04}$	$9.39^{+2.49}_{-3.88}$	5.61 ± 0.06
A3526	$3.69^{+0.05}_{-0.04}$	1.19 ± 0.04	$3.41^{+0.60}_{-1.36}$	$1.32^{+0.07}_{-0.20}$	$0.11^{+0.04}_{-0.01}$	$3.83^{+0.39}_{-0.75}$	3.66 ± 0.16
A3558	$5.37^{+0.17}_{-0.15}$	6.56 ± 0.04	$6.71^{+0.91}_{-2.12}$	$1.59^{+0.07}_{-0.19}$	$0.22^{+0.05}_{-0.02}$	$14.60^{+0.84}_{-2.35}$	11.10 ± 0.06
A3562	$4.47^{+0.23}_{-0.21}$	3.08 ± 0.05	$3.51^{+0.43}_{-0.91}$	$1.28^{+0.05}_{-0.12}$	$0.20^{+0.04}_{-0.02}$	$7.07^{+0.50}_{-1.10}$	3.23 ± 0.08
A3571	$6.80^{+0.21}_{-0.18}$	8.08 ± 0.11	$8.76^{+1.69}_{-3.43}$	$1.75^{+0.10}_{-0.27}$	$0.19^{+0.07}_{-0.02}$	$16.45^{+1.36}_{-2.93}$	-
A3581	$1.83^{+0.04}_{-0.02}$	0.60 ± 0.03	$0.93^{+0.38}_{-0.44}$	$0.84^{+0.14}_{-0.05}$	$0.15^{+0.03}_{-0.01}$	$1.41^{+0.29}_{-0.41}$	1.22 ± 0.09
A3667	$6.28^{+0.27}_{-0.26}$	9.48 ± 0.11	$5.28^{+0.52}_{-1.15}$	$1.46^{+0.05}_{-0.11}$	$0.30^{+0.03}_{-0.01}$	$16.04^{+1.15}_{-2.22}$	8.65 ± 0.06
A4038	$3.22^{+0.10}_{-0.10}$	1.92 ± 0.04	$2.58^{+0.49}_{-1.05}$	$1.18^{+0.07}_{-0.19}$	$0.16^{+0.08}_{-0.03}$	$4.14^{+0.61}_{-0.92}$	2.85 ± 0.06
A4059	$3.94^{+0.15}_{-0.15}$	2.80 ± 0.06	$4.41^{+1.14}_{-2.03}$	$1.39^{+0.11}_{-0.26}$	$0.14^{+0.08}_{-0.02}$	$6.17^{+0.79}_{-1.40}$	3.12 ± 0.10
COMA	$8.07^{+0.29}_{-0.27}$	8.09 ± 0.19	$9.95^{+2.10}_{-2.99}$	$1.86^{+0.12}_{-0.21}$	$0.19^{+0.07}_{-0.04}$	$19.00^{+1.89}_{-3.66}$	8.94 ± 0.05
EXO0422	$2.90^{+0.90}_{-0.60}$	2.03 ± 0.21	$2.72^{+1.71}_{-1.45}$	$1.19^{+0.21}_{-0.27}$	$0.14^{+0.13}_{-0.06}$	$3.68^{+2.04}_{-1.75}$	1.63 ± 0.08
FORNAX	$1.56^{+0.05}_{-0.07}$	0.08 ± 0.01	$1.29^{+0.55}_{-0.44}$	$0.96^{+0.16}_{-0.16}$	$0.04^{+0.01}_{-0.01}$	$0.46^{+0.14}_{-0.14}$	-
HYDRA-A	$3.82^{+0.20}_{-0.17}$	5.84 ± 0.04	$4.07^{+1.27}_{-2.14}$	$1.34^{+0.13}_{-0.29}$	$0.18^{+0.11}_{-0.03}$	$7.29^{+0.77}_{-1.72}$	3.09 ± 0.02
IIIZw54	$(3.00^{+1.39}_{-0.95})$	0.83 ± 0.10	$3.76^{+2.82}_{-2.26}$	$1.33^{+0.27}_{-0.35}$	$0.08^{+0.12}_{-0.05}$	$2.95^{+1.97}_{-1.72}$	-
MKW3S	$3.45^{+0.13}_{-0.10}$	2.79 ± 0.05	$3.22^{+0.92}_{-1.53}$	$1.25^{+0.11}_{-0.24}$	$0.15^{+0.08}_{-0.02}$	$4.92^{+0.59}_{-1.06}$	1.96 ± 0.08
MKW4	$1.84^{+0.05}_{-0.03}$	0.34 ± 0.01	$0.69^{+0.04}_{-0.14}$	$0.76^{+0.01}_{-0.06}$	$0.15^{+0.02}_{-0.01}$	$1.05^{+0.07}_{-0.15}$	1.71 ± 0.23
MKW8	$3.29^{+0.23}_{-0.22}$	0.79 ± 0.11	$2.00^{+0.46}_{-0.74}$	$1.08^{+0.08}_{-0.12}$	$0.11^{+0.08}_{-0.01}$	$2.22^{+1.11}_{-1.29}$	2.13 ± 0.08
NGC1550	$1.44^{+0.03}_{-0.02}$	0.28 ± 0.02	$0.68^{+0.14}_{-0.24}$	$0.77^{+0.05}_{-0.10}$	$0.12^{+0.07}_{-0.04}$	$0.79^{+0.29}_{-0.33}$	-
NGC4636	$0.66^{+0.03}_{-0.01}$	0.02 ± 0.00	$0.18^{+0.03}_{-0.06}$	$0.49^{+0.03}_{-0.07}$	$0.04^{+0.03}_{-0.02}$	$0.08^{+0.04}_{-0.04}$	-
NGC5044	$1.22^{+0.04}_{-0.04}$	0.18 ± 0.00	$0.49^{+0.12}_{-0.25}$	$0.69^{+0.05}_{-0.15}$	$0.09^{+0.04}_{-0.01}$	$0.45^{+0.05}_{-0.14}$	-
NGC507	$1.40^{+0.04}_{-0.07}$	0.23 ± 0.00	$0.46^{+0.02}_{-0.07}$	$0.67^{+0.01}_{-0.04}$	$0.15^{+0.02}_{-0.01}$	$0.68^{+0.04}_{-0.08}$	1.94 ± 0.16
S1101	$2.60^{+0.50}_{-0.50}$	3.52 ± 0.05	$2.94^{+1.57}_{-1.88}$	$1.20^{+0.18}_{-0.34}$	$0.14^{+0.15}_{-0.04}$	$4.22^{+0.90}_{-1.23}$	2.18 ± 0.08
ZwCl1215	$(6.36^{+2.94}_{-2.01})$	5.17 ± 0.11	$9.46^{+3.74}_{-4.87}$	$1.74^{+0.30}_{-0.37}$	$0.15^{+0.08}_{-0.04}$	$13.87^{+3.01}_{-3.76}$	-

Note: The values of T_h and L_X are from Ikebe et al. (2002). T_h in a bracket is estimated from the $L_X - T$ relation (Ikebe et al. 2002). T_m with ‘a’ is derived from non-ASCA spectroscopy (Ikebe et al. 2002 and references therein).

Table 5. Continued (extended cluster sample).

Name	T_h (keV)	L_X (10^{44} erg/s, 0.1-2.4keV)	M_{500} ($10^{14} M_\odot$)	r_{500} (Mpc)	f_{gas}	$M_{gas,500}$ ($10^{13} M_\odot$)	L_{mir} ($h_{70}^{-2} 10^{12} L_\odot$)
3C129	$5.57^{+0.16}_{-0.15}$	2.27 ± 0.21	$5.39^{+2.26}_{-2.33}$	$1.51^{+0.19}_{-0.26}$	$0.16^{+0.16}_{-0.10}$	$8.55^{+2.58}_{-4.82}$	-
A0539	$3.04^{+0.10}_{-0.10}$	1.11 ± 0.02	$2.68^{+0.32}_{-0.85}$	$1.19^{+0.65}_{-0.14}$	$0.14^{+0.05}_{-0.01}$	$3.78^{+0.27}_{-0.47}$	3.72 ± 0.05
A0548e	$2.93^{+0.17}_{-0.15}$	1.05 ± 0.03	$1.49^{+0.85}_{-0.17}$	$0.97^{+0.03}_{-0.04}$	$0.18^{+0.02}_{-0.02}$	$2.65^{+0.27}_{-0.29}$	4.17 ± 0.08
A0548w	$(1.68^{+0.77}_{-0.53})$	0.19 ± 0.02	$1.00^{+0.58}_{-0.52}$	$0.85^{+0.14}_{-0.19}$	$0.08^{+0.07}_{-0.04}$	$0.82^{+0.45}_{-0.43}$	-
A0644	$6.54^{+0.27}_{-0.26}$	8.35 ± 0.15	$8.41^{+2.15}_{-3.81}$	$1.68^{+0.13}_{-0.31}$	$0.16^{+0.07}_{-0.02}$	$13.85^{+1.56}_{-2.97}$	6.34 ± 0.06
A1413	$6.56^{+0.65}_{-0.44}$	10.71 ± 0.30	$9.77^{+2.78}_{-4.58}$	$1.65^{+0.14}_{-0.31}$	$0.17^{+0.09}_{-0.03}$	$16.85^{+1.94}_{-3.46}$	-
A1689	$8.58^{+0.84}_{-0.40}$	19.48 ± 0.34	$14.98^{+5.82}_{-8.38}$	$1.84^{+0.21}_{-0.44}$	$0.16^{+0.13}_{-0.03}$	$23.26^{+3.69}_{-5.42}$	-
A1775	$3.66^{+0.34}_{-0.25}$	3.09 ± 0.09	$4.19^{+1.36}_{-1.69}$	$1.32^{+0.13}_{-0.21}$	$0.18^{+0.07}_{-0.03}$	$7.55^{+2.21}_{-2.07}$	-
A1800	$(5.02^{+2.32}_{-1.59})$	2.85 ± 0.37	$5.94^{+4.83}_{-3.59}$	$1.49^{+0.33}_{-0.40}$	$0.13^{+0.16}_{-0.09}$	$7.63^{+4.25}_{-4.68}$	-
A1914	$8.41^{+0.60}_{-0.58}$	17.04 ± 0.38	$11.84^{+3.65}_{-5.84}$	$1.72^{+0.16}_{-0.35}$	$0.18^{+0.10}_{-0.03}$	$21.20^{+2.54}_{-4.72}$	-
A2151w	$2.58^{+0.19}_{-0.20}$	0.89 ± 0.03	$1.60^{+0.35}_{-0.61}$	$1.00^{+0.07}_{-0.15}$	$0.14^{+0.05}_{-0.02}$	$2.25^{+0.24}_{-0.51}$	3.44 ± 0.14
A2319	$8.84^{+0.29}_{-0.24}$	16.37 ± 0.26	$13.57^{+2.15}_{-4.59}$	$1.99^{+0.10}_{-0.26}$	$0.24^{+0.07}_{-0.02}$	$32.26^{+3.88}_{-5.92}$	14.63 ± 0.04
A2734	$5.07^{+0.36}_{-0.42}$	2.40 ± 0.10	$4.82^{+0.85}_{-1.56}$	$1.40^{+0.68}_{-0.17}$	$0.14^{+0.05}_{-0.02}$	$6.80^{+0.97}_{-1.57}$	3.13 ± 0.09
A2877	$^a 3.50^{+2.20}_{-1.10}$	0.40 ± 0.01	$6.88^{+6.74}_{-3.79}$	$1.64^{+0.42}_{-0.38}$	$0.04^{+0.03}_{-0.02}$	$3.04^{+0.51}_{-0.63}$	-
A3395n	$5.11^{+0.47}_{-0.43}$	1.63 ± 0.11	$8.13^{+5.53}_{-4.69}$	$1.69^{+0.32}_{-0.42}$	$0.10^{+0.12}_{-0.07}$	$8.24^{+2.20}_{-5.34}$	-
A3528n	$4.79^{+0.50}_{-0.44}$	1.56 ± 0.06	$4.49^{+0.78}_{-1.53}$	$1.38^{+0.08}_{-0.18}$	$0.12^{+0.04}_{-0.02}$	$5.33^{+0.67}_{-1.18}$	-
A3528s	$4.60^{+0.49}_{-0.27}$	2.20 ± 0.06	$2.76^{+0.39}_{-0.44}$	$1.17^{+0.05}_{-0.07}$	$0.20^{+0.03}_{-0.02}$	$5.56^{+0.51}_{-0.73}$	-
A3530	$4.05^{+0.32}_{-0.30}$	1.21 ± 0.06	$4.34^{+1.09}_{-1.27}$	$1.37^{+0.11}_{-0.15}$	$0.12^{+0.05}_{-0.04}$	$5.22^{+0.89}_{-1.38}$	-
A3532	$4.41^{+0.19}_{-0.18}$	2.20 ± 0.06	$6.63^{+1.17}_{-1.84}$	$1.57^{+0.09}_{-0.27}$	$0.12^{+0.08}_{-0.01}$	$8.11^{+0.68}_{-1.32}$	-
A3560	$(3.90^{+1.81}_{-1.23})$	1.57 ± 0.06	$2.77^{+1.85}_{-1.26}$	$1.18^{+0.32}_{-0.22}$	$0.17^{+0.06}_{-0.04}$	$4.61^{+1.32}_{-1.36}$	6.50 ± 0.27
A3627	$5.62^{+0.12}_{-0.11}$	3.59 ± 0.18	$4.92^{+0.67}_{-0.91}$	$1.48^{+0.06}_{-0.10}$	$0.19^{+0.05}_{-0.04}$	$9.57^{+1.37}_{-2.14}$	-
A3695	$(6.76^{+3.12}_{-2.14})$	5.89 ± 0.89	$7.03^{+4.66}_{-4.22}$	$1.55^{+0.29}_{-0.41}$	$0.18^{+0.18}_{-0.12}$	$12.82^{+6.66}_{-7.84}$	-
A3822	$5.12^{+0.43}_{-0.31}$	4.83 ± 0.58	$4.69^{+1.19}_{-1.46}$	$1.37^{+0.11}_{-0.16}$	$0.19^{+0.13}_{-0.08}$	$9.07^{+2.53}_{-4.06}$	5.38 ± 0.10
A3827	$(7.66^{+3.54}_{-2.42})$	7.94 ± 0.78	$15.50^{+3.81}_{-9.76}$	$2.01^{+0.48}_{-0.56}$	$0.13^{+0.19}_{-0.08}$	$19.88^{+11.11}_{-11.45}$	-
A3888	$(8.68^{+4.01}_{-2.75})$	10.09 ± 0.39	$29.81^{+21.73}_{-1.50}$	$2.38^{+0.54}_{-0.84}$	$0.07^{+0.13}_{-0.03}$	$21.97^{+3.60}_{-6.63}$	-
A3921	$5.39^{+0.38}_{-0.35}$	4.83 ± 0.14	$6.59^{+1.50}_{-2.32}$	$1.51^{+0.11}_{-0.20}$	$0.16^{+0.06}_{-0.03}$	$10.76^{+1.41}_{-2.05}$	-
AWM7	$3.70^{+0.08}_{-0.05}$	2.10 ± 0.07	$4.92^{+1.21}_{-2.26}$	$1.48^{+0.11}_{-0.27}$	$0.12^{+0.07}_{-0.02}$	$5.85^{+0.70}_{-1.10}$	2.90 ± 0.06
HCG94	$3.30^{+0.17}_{-0.16}$	1.28 ± 0.02	$2.25^{+0.24}_{-0.60}$	$1.11^{+0.04}_{-0.11}$	$0.15^{+0.03}_{-0.01}$	$3.41^{+0.18}_{-0.49}$	2.51 ± 0.11
II Zw108	$(4.28^{+1.98}_{-1.35})$	1.98 ± 0.24	$3.85^{+2.60}_{-2.00}$	$1.32^{+0.25}_{-0.28}$	$0.15^{+0.11}_{-0.07}$	$5.65^{+2.88}_{-2.73}$	-
M49	$1.33^{+0.03}_{-0.02}$	0.02 ± 0.00	$0.67^{+0.40}_{-0.47}$	$0.77^{+0.13}_{-0.26}$	$0.01^{+0.01}_{-0.00}$	$0.09^{+0.02}_{-0.04}$	-
NGC499	$0.66^{+0.03}_{-0.03}$	0.04 ± 0.00	$0.33^{+0.25}_{-0.23}$	$0.60^{+0.20}_{-0.12}$	$0.03^{+0.04}_{-0.01}$	$0.08^{+0.03}_{-0.03}$	-
NGC5813	$0.76^{+0.19}_{-0.19}$	0.02 ± 0.00	$0.43^{+0.45}_{-0.33}$	$0.66^{+0.18}_{-0.26}$	$0.01^{+0.05}_{-0.01}$	$0.06^{+0.07}_{-0.04}$	-
NGC5846	$0.64^{+0.04}_{-0.03}$	0.01 ± 0.00	$0.18^{+0.11}_{-0.12}$	$0.49^{+0.08}_{-0.16}$	$0.03^{+0.03}_{-0.01}$	$0.05^{+0.02}_{-0.02}$	-
OPHIUCHU	$10.25^{+0.30}_{-0.36}$	12.14 ± 0.39	$38.76^{+21.59}_{-24.11}$	$2.91^{+0.46}_{-0.81}$	$0.07^{+0.09}_{-0.02}$	$28.60^{+2.89}_{-6.11}$	-
PERSEUS	$6.42^{+0.06}_{-0.06}$	16.39 ± 0.20	$6.08^{+1.55}_{-2.85}$	$1.58^{+0.12}_{-0.30}$	$0.30^{+0.12}_{-0.03}$	$18.10^{+1.81}_{-4.79}$	8.63 ± 0.07
PKS0745	$6.37^{+0.21}_{-0.20}$	27.13 ± 0.42	$7.11^{+2.85}_{-3.91}$	$1.54^{+0.18}_{-0.36}$	$0.28^{+0.18}_{-0.05}$	$19.74^{+2.46}_{-5.35}$	-
RXJ2344	$(5.52^{+2.55}_{-1.74})$	3.59 ± 0.08	$8.89^{+5.48}_{-5.25}$	$1.70^{+0.29}_{-0.44}$	$0.11^{+0.10}_{-0.03}$	$9.47^{+1.70}_{-2.29}$	-
S405	$(5.02^{+2.32}_{-1.59})$	2.90 ± 0.39	$4.62^{+3.14}_{-2.88}$	$1.39^{+0.26}_{-0.39}$	$0.16^{+0.13}_{-0.10}$	$7.59^{+3.71}_{-4.37}$	-
S540	$(3.09^{+1.43}_{-0.98})$	0.89 ± 0.07	$2.52^{+1.55}_{-1.40}$	$1.16^{+0.20}_{-0.27}$	$0.11^{+0.08}_{-0.05}$	$2.68^{+1.18}_{-1.23}$	-
S636	$9.06^{+0.07}_{-0.06}$	0.38 ± 0.03	$1.56^{+0.44}_{-0.47}$	$1.01^{+0.09}_{-0.11}$	$0.10^{+0.07}_{-0.04}$	$1.52^{+0.35}_{-0.62}$	-
TRIANGUL	$2.06^{+0.33}_{-0.31}$	12.43 ± 0.15	$14.84^{+2.49}_{-5.28}$	$2.07^{+0.11}_{-0.28}$	$0.20^{+0.06}_{-0.02}$	$29.14^{+2.35}_{-4.92}$	12.01 ± 0.05
UGC03957	$(3.21^{+1.48}_{-1.02})$	0.98 ± 0.10	$3.32^{+2.76}_{-2.03}$	$1.28^{+0.34}_{-0.34}$	$0.08^{+0.10}_{-0.04}$	$2.67^{+1.62}_{-1.47}$	-
ZwCl1742	$(6.05^{+2.80}_{-1.91})$	4.54 ± 0.33	$10.11^{+5.91}_{-6.30}$	$1.78^{+0.30}_{-0.49}$	$0.11^{+0.13}_{-0.03}$	$11.20^{+3.42}_{-3.22}$	-

Note: The symbols in T_h have the same meanings as that in Table 4.

L_X - M relation is again higher than 1.0 which is predicted by a self-similar model, as shown in Fig. 8.

Like in the X-ray luminosity temperature relation we see a substantial difference in the normalization of this relation for the CCCs and NCCCs by about a factor of 2.4. The offset between the different cluster subsamples shows that the scatter in the overall sample is partly produced by the different types of clusters and knowing more about the clusters, this scatter could be reduced as discussed below.

One of the important questions concerns the origin of this large difference. There are in principle two effects

caused by cool cores which add to the observed result: cluster temperatures and the cluster masses derived from temperature estimates will be biased low and luminosities will be biased high due to the enhanced emission of dense cores and more compact clusters. In turn if the core radii of the NCCCs are inflated, the cluster masses will be biased high for the NCCCs.

To distinguish these different possibilities we need mass estimators independent of the temperature and independent of the core radii. In the next section we apply such new parameters.




Fig. 6. L_X - T_m relation. The solid line shows the power law fit to all the data, and the dotted and dashed lines are those for the CCCs and NCCCs, respectively. T_m with a, b, c and d in Table 1 are selected to plot here. Note T_m with d is replaced by the temperature measured from the central 2' or 3' region (see Table 2 in Fukazawa et al. 2000).




Fig. 8. L_X - M_{500} relation. The dot-dashed line shows the power law relation fit to all the data (106 clusters). The solid line shows that for all the data with ASCA measured T_h , and the dotted and dashed lines are those for the CCCs and NCCCs, respectively.




Fig. 7. L_X - T_h relation. The solid line shows the power law relation fit to all the data, and the dotted and dashed lines are those for the CCCs and NCCCs, respectively.

Fig. 8 also shows that the fraction of CCCs at the high L_X end do not constitute a small fraction of all clusters unlike that at the high M_{500} end. This is due to the fact that CCCs usually have higher L_X for the clusters with the same M_{500} . The CCC fraction is about 60% (11 CCCs) for the most luminous 18 clusters (with $L_X > 6 \cdot 10^{44}$ erg/s and $z < 0.17$ in the Λ CDM cosmology) in this sample. This fraction is consistent with that in the distant cluster samples with similar L_X , e.g., 7 CCCs out of 12 clusters at $z \sim 0.2$ (Zhang et al. 2006b), 6 CCCs out of 13 clusters




Fig. 9. L_{nir} - T_h relation. The solid line shows the power law relation fit to all the data and the dotted and dashed lines are those for the CCCs and NCCCs, respectively.

at $z \sim 0.3$ (Zhang et al. 2006a) where we use the same criterion of CCCs as in the HIFLUCS. This result shows that the fraction of CCCs in luminous cluster samples does not show a large evolutionary effect up to $z \sim 0.3$.

4.5. Relations involving total NIR luminosity and gas mass

Assuming that the cluster gas mass fraction is approximately constant with cluster mass (e.g. Allen et al. 2004, Ettori et al. 2003), we can also use the cluster gas mass




Fig. 10. $L_{nir} - M_{gas,500}$ relation. The solid line shows the power law relation fit to all the data, and the dotted and dashed lines are those for the CCCs and NCCCs, respectively. Note that NCCCs have larger near infrared luminosities L_{nir} than CCCs.




Fig. 12. $M_{gas,500} - T_m$ relation. T_m with a, b, c and d in Table 1 are selected to plot here. Note T_m with d is replaced by the temperature measured from the central 2' or 3' region (see Table 2 in Fukazawa et al. 2000). Symbols have the same meanings as in Fig. 11.




Fig. 11. $M_{gas,500} - T_h$ relation. The solid line shows the power law relation fit to all the data, and the dotted and dashed lines are those for the CCCs and NCCCs, respectively.

as an estimator of the cluster's total mass. The determination of the cluster gas mass depends only on the X-ray surface brightness distribution, not on the temperature. However, the way we determine $M_{gas,500}$, based on the fiducial outer radius of r_{500} , introduces a weak temperature dependence, since in our approach we used the temperature dependent gravitational mass to estimate r_{500} . This dependence is roughly proportional to $T^{1/2}$ and thus much weaker than proportional and much weaker than the dependence of $M_{500} \propto T^{3/2}$. Thus if there is any strong bias in the temperature determination due to cooling flows




Fig. 13. $f_{gas,500} - T_h$ relation.

we should still see this effect in a correlation analysis based on gas mass, but it will be just weakened approximately by a factor of two. For example, if the temperature of a cluster, T , is biased to $2T$, the measured $M_{gas,500}$ will change to $1.4M_{gas,500}$. However, if this cluster remains on the line with a slope 2.0 (see Table 7), it need that $M_{gas,500}$ change to $4M_{gas,500}$. From this example, one can see that $M_{gas,500}$ is insensitive to T in the relation $M_{gas,500} - T$. Another mass estimator is the total luminosity of the cluster galaxies in the NIR (K-band), that is obtained from the Two Micron All Sky Survey (2MASS) (Lin et al. 2004).

The $L_{nir} - T_h$ relation in Fig. 9 shows no strong bias of CCCs versus NCCCs. The normalization difference is less

than 10% and well within the 1σ error of the fits. The comparison of the two mass estimators L_{nir} and $M_{gas,500}$ in Fig. 10 shows a comparatively small difference between the CCCs and NCCCs with a factor smaller than about 1.25, where the CCCs have on average a little higher $M_{gas,500}$. Therefore we do not see a strong bias in either one of the two mass estimators, although a weak bias cannot be ruled out, especially in the gas mass.

Similarly, in Figs. 11 and 12, where we show the $M_{gas,500}-T_h$ and $M_{gas,500}-T_m$ relations, we do not see a strong temperature bias for CCCs versus NCCCs. Similar to Figs. 6 and 7, the slope of $M_{gas,500}-T_m$ is shallower than that of $M_{gas,500}-T_h$, but it only has a slight difference for the clusters with $T_h > 3\text{keV}$. Note that clusters with $T_m < 1.0\text{keV}$ in Fig. 11 are not plotted in Fig. 12. In addition, the difference in the normalization of this relation for the CCCs and NCCCs is very small and within the measurement errors, with a factor 1.10 and 1.00 for T_m and T_h , respectively. Therefore we conclude from the results in this section, that the segregation of CCCs and NCCCs in the L_X-M and L_X-T relations is to the major part an X-ray luminosity effect and to a lesser extent an effect of a biased temperature estimate. A similar conclusion has been reached by O'Hara et al. (2006).

4.6. Gas mass fraction

We found no difference of the total gas mass fraction, $f_{gas,500}$, between CCCs and NCCCs as shown in Fig. 13. This reconfirms the small influence of cooling cores on the mass and gas mass estimates.

5. Discussions and summary

In this paper we use an isothermal model to determine the X-ray mass, due to the fact that only global temperature estimates mainly from ASCA are available for such a large sample up to date. For the central region in cooling core clusters this is obviously not correct. But at large radii we expect that the total gravitational mass is correct. This is confirmed by our mass determination for the cluster PKS 0745-191 and Abell 1650 based on XMM-Newton observations (Chen et al. 2003, Jia et al. 2006) in which the resultant total masses are found to be consistent with this work.

In summary we find from the analysis presented in this paper that:

(i) the formally derived mass deposition rates for the strongest cooling core clusters are roughly proportional to the cluster mass.

(ii) the fraction of NCCCs increases significantly with M_{500} and most of the galaxy groups in HIFLUGCS are cooling core clusters. This is most probably explained by the fact that the most massive galaxy clusters have been formed most recently and should therefore show a larger fraction of dynamically young systems. These may turn into cooling core clusters in a later evolutionary stage. In

addition, the fraction of CCCs in luminous cluster samples does not show a large evolution effect within $z < 0.3$.

(iii) among all the observational parameters, the core radius and the X-ray luminosity are shown to be most affected by the presence of a cooling core, as observed in their relation to other bulk cluster properties as cluster mass and temperature.

(iv) the $M-T$ relation using the X-ray temperature of the hot ICM phase, T_h , seems to show a comparatively small bias for CCCs in comparison to NCCCs.

From the magnitude of the effect (iii) we can conclude that the scatter in the $L_X - M_{500}$ relation, which is so important in cosmological applications, is to a large part due to the different normalizations of CCCs and NCCCs. It is important to distinguish between statistical, systematic, and intrinsic scatter and to take the intrinsic scatter into account in applications (e.g. Ikebe et al. 2002; Stanek et al. 2006). The findings here indicate that a significant portion of the scatter may be intrinsic due to variations in the X-ray luminosity for clusters of a given mass. We are currently working to confirm this result with high quality Chandra and XMM-Newton observations of the complete HIFLUGCS sample. If a substantial scatter is confirmed, it will be interesting to check if these results are fully consistent with the low Ω_m and σ_8 values found from the WMAP 3rd year data (Spergel et al. 2006) combined with cluster mass function prediction and observed luminosity function (Reiprich 2006). This finding is also pointing the way to an improvement of the $L_X - M_{500}$ relation. Given a good proxy for the strength of cooling cores, one could correct for this effect and tighten this important relation. This was indeed suggested by O'Hara et al. (2006) who propose to use the central surface brightness as such a proxy. Further work is in progress to use the HIFLUGCS cluster sample to work out such a correction scheme.

Acknowledgements. Y. Chen was supported by the exchange program between the Max-Planck Society and the Chinese Academy of Sciences. He thanks MPE for its very gracious hospitality. This research is partially supported by the Scientific Research Foundation for the Returned Overseas Chinese Scholars, State Education Ministry.

References

- Akritas, M.G. & Bershady, M.A. 1996, ApJ, 470, 706
- Allen, S.W., Schmidt, R.W., Ebeling, H., Fabian, A.C., van Speybroeck, L., 2004, MNRAS, 353, 457
- Arnaud, M., Pointecouteau, E., Pratt, G.W., 2005, A&A, 441, 893
- Arnaud, K.A. 1996, ADASS, 101, 5
- Böhringer, H., Matsushita, K., Churazov, E., Ikebe, Y. & Chen, Y. 2002, A&A, 382, 804
- Cavaliere, A. & Fusco-Femiano, R. 1976, A&A, 49, 137
- Chen, Y., Ikebe, Y. & Böhringer, H., 2003, A&A, 407, 41
- David, L.P., Nulsen, P.E.J., McNamara, B.R., 2001, ApJ, 557, 546
- Edge, A.C. & Stewart, G.C., 1991, MNRAS, 252, 414
- Ettori, S. 2000, MNRAS, 311, 313
- Ettori, S., De Grandi, S. & Molendi, S. 2002, A&A, 391, 841

Table 6. Fits to relations of r_c and M_{500} . The relations are given in the form: $\log_{10} \left(\frac{r_c}{100 \text{ kpc}} \right) = A + B \cdot \log_{10} \left(\frac{M_{500}}{5 \cdot 10^{14} M_{\odot}} \right)$.

relation	Number of clusters	B	A	comments
$r_c - M_{500}$	106	1.18 ± 0.08	0.253 ± 0.036	ALL
$r_c - M_{500}$	52	0.90 ± 0.06	0.023 ± 0.036	CCCs
$r_c - M_{500}$	54	0.66 ± 0.12	0.452 ± 0.029	NCCCs

Table 7. Summary of the fits to the scaling relations. The relations are given in the form: $\log_{10} \left(\frac{M_{500}}{5 \cdot 10^{14} M_{\odot}} \right) = A + B \cdot \log_{10} \left(\frac{T}{4 \text{ keV}} \right)$, $\log_{10} \left(\frac{M_{gas,500}}{10^{14} M_{\odot}} \right) = A + B \cdot \log_{10} \left(\frac{T}{4 \text{ keV}} \right)$, $\log_{10} \left(\frac{L_X}{10^{44} \text{ erg/s}} \right) = A + B \cdot \log_{10} \left(\frac{T}{4 \text{ keV}} \right)$, $\log_{10} \left(\frac{L_X}{10^{44} \text{ erg/s}} \right) = A + B \cdot \log_{10} \left(\frac{M_{500}}{5 \cdot 10^{14} M_{\odot}} \right)$, $\log_{10} (L_{nir}) = A + B \cdot \log_{10} \left(\frac{T}{4 \text{ keV}} \right)$ and $\log_{10} (L_{nir}) = A + B \cdot \log_{10} \left(\frac{M_{gas,500}}{10^{14} M_{\odot}} \right)$.

relation	Number of clusters	B	A	comments
$M_{500}-T_h$	88	1.54 ± 0.06	-0.112 ± 0.014	ALL with T_h
$M_{500}-T_h$	47	1.48 ± 0.07	-0.140 ± 0.015	CCCs
$M_{500}-T_h$	41	1.57 ± 0.17	-0.088 ± 0.031	NCCCs
$M_{500}-T_h$	72	1.30 ± 0.18	-0.069 ± 0.025	$T_h > 3.0 \text{ keV}$
$M_{500}-T_h$	16	0.91 ± 0.31	-0.46 ± 0.13	$T_h < 3.0 \text{ keV}$
$M_{gas,500}-T_h$	88	2.29 ± 0.09	-0.269 ± 0.015	ALL with T_h
$M_{gas,500}-T_h$	47	2.38 ± 0.10	-0.251 ± 0.017	CCCs
$M_{gas,500}-T_h$	41	2.04 ± 0.14	-0.258 ± 0.025	NCCCs
$M_{gas,500}-T_h$	72	1.89 ± 0.10	-0.221 ± 0.017	$T_h > 3.0 \text{ keV}$
$M_{gas,500}-T_h$	35	1.98 ± 0.11	-0.219 ± 0.019	$T_h > 3.0 \text{ keV}$, CCCs
$M_{gas,500}-T_h$	37	1.80 ± 0.18	-0.219 ± 0.033	$T_h > 3.0 \text{ keV}$, NCCCs
$M_{gas,500}-T_m$	71	1.70 ± 0.06	-0.184 ± 0.015	ALL with T_m
$M_{gas,500}-T_m$	38	1.73 ± 0.07	-0.166 ± 0.022	CCCs
$M_{gas,500}-T_m$	33	1.71 ± 0.11	-0.207 ± 0.025	NCCCs
L_X-T_h	88	2.73 ± 0.13	0.363 ± 0.027	ALL with T_h
L_X-T_h	47	2.88 ± 0.15	0.492 ± 0.031	CCCs
L_X-T_h	41	2.74 ± 0.17	0.227 ± 0.034	NCCCs
L_X-T_h	72	2.88 ± 0.19	0.332 ± 0.034	$T_h > 3.0 \text{ keV}$
L_X-T_h	35	3.08 ± 0.24	0.457 ± 0.039	$T_h > 3.0 \text{ keV}$, CCCs
L_X-T_h	37	2.78 ± 0.22	0.213 ± 0.044	$T_h > 3.0 \text{ keV}$, NCCCs
L_X-T_m	71	2.23 ± 0.11	0.458 ± 0.032	ALL with T_m
L_X-T_m	38	2.31 ± 0.14	0.597 ± 0.040	CCCs
L_X-T_m	33	2.33 ± 0.09	0.286 ± 0.031	NCCCs
L_X-M_{500}	106	1.82 ± 0.13	0.521 ± 0.039	ALL
L_X-M_{500}	88	1.77 ± 0.12	0.562 ± 0.041	ALL with T_h
L_X-M_{500}	47	1.94 ± 0.15	0.763 ± 0.050	CCCs
L_X-M_{500}	41	1.75 ± 0.25	0.381 ± 0.062	NCCCs
L_X-M_{500}	72	2.23 ± 0.38	0.485 ± 0.053	$T_h > 3.0 \text{ keV}$
L_X-M_{500}	35	3.05 ± 0.60	0.674 ± 0.074	$T_h > 3.0 \text{ keV}$, CCCs
L_X-M_{500}	37	1.84 ± 0.42	0.350 ± 0.087	$T_h > 3.0 \text{ keV}$, NCCCs
$L_{nir}-T_h$	58	1.34 ± 0.09	0.586 ± 0.017	ALL with T_m
$L_{nir}-T_h$	31	1.17 ± 0.11	0.565 ± 0.017	CCCs
$L_{nir}-T_h$	27	1.47 ± 0.15	0.593 ± 0.035	NCCCs
$L_{nir}-M_{gas,500}$	62	0.702 ± 0.055	0.741 ± 0.016	ALL
$L_{nir}-M_{gas,500}$	33	0.604 ± 0.062	0.686 ± 0.018	CCCs
$L_{nir}-M_{gas,500}$	29	0.756 ± 0.080	0.778 ± 0.024	NCCCs

Ettori, S., Tozzi, P., Rosati, P., 2003, A&A, 398, 879

Fabian, A.C., 1994, ARA&A, 32, 277

Fabian, A.C., Crawford, C.S., Edge, A.C., Mushotzky, R.F., 1994, MNRAS, 267, 779

Fabian, A.C., Sanders, J.S., Allen, S.W., et al., 2003, MNRAS, 344, L43

Finoguenov, A., Reiprich, T.H. & Böhringer, H., 2001, A&A, 368, 749

Fukazawa, Y., Makishima, K., Tamura, T., et al., 1998, PASJ, 50, 187

Fukazawa, Y., Makishima, K., Tamura, T., et al., 2000, MNRAS, 313, 21

Ikebe, Y., Makishima, K., Fukazawa, Y., et al., 1999, ApJ, 525, 58

Ikebe, Y., Reiprich, T.H., Böhringer, H. et al. 2002, A&A, 383, 773

- Jia, S.M., Chen, Y. & Chen, L., 2006, ChJAA, Vol. 6, 181
- Jones, C., Forman, W., 1984, ApJ, 276, 38
- Lin, Y.T., Mohr, J.J. & Stanford, S.A., 2004, ApJ, 617, 879
- Makishima, K., Ezawa, H., Fukuzawa, Y. et al., 2001, PASJ, 53, 401
- Markevitch, M., Forman, W.R., Sarazin, C.L., Vikhlinin, A., 1998, ApJ, 503, 77
- Markevitch, M., 1998, ApJ, 504, 27
- McCarthy, I.G., Balogh, M.L., Babul, A., Poole, G.B., Horner, D.J., 2004, ApJ, 613, 811
- McNamara, B.R., Nulsen, P.E.J., Wise, M.W., et al., 2005, Nature, 433, 45
- Molendi, S. & Pizzolato, F. 2001, ApJ, 560, 194
- O'Hara, T.B., Mohr, J.J., Bialek, J.J., Evrard, A.E., 2006, ApJ, 639, 640
- Ota, N. & Mitsuda, K., 2004, A&A, 428, 757
- Peres, C.B., Fabian, A.C., Edge, A.C., et al. 1998, MNRAS, 298, 416
- Peterson, J.R., Paerels, F.B.S., Kaastra, J.S., et al., 2001, A&A, 365, L104
- Peterson, J.R., Kahn, S.M., Paerels, F.B.S., et al., 2003, ApJ, 590, 207
- Pratt, G.W., et al. 2006, A&A, submitted
- Reiprich, T.H. 2001, Cosmological Implications and Physical Properties of an X-ray Flux-Limited Sample of Galaxy Clusters, Ph.D. thesis
- Reiprich, T.H. & Böhringer H. 2002, ApJ, 567, 716
- Reiprich, T.H., 2006, A&A, 453, L39
- Schuecker, P., Böhringer, H., Reiprich, T.H., Feretti, L., 2001a, A&A, 378, 408
- Schuecker, P., Böhringer, H., Guzzo, L., Collins, C.A., et al., 2001b, A&A, 368, 86
- Schuecker, P., Böhringer, H., Collins, C.A., Guzzo, L., 2003, A&A, 398, 867
- Spergel, D.N., Beam, R., Doré, O., et al., 2006, ApJ, submitted, [astro-ph/0603449]
- Stanek, R., Evrard, A.E., Böhringer, H., Schuecker, P., Nord, B., 2006, astro-ph/0602324
- Vikhlinin, A., Markevitch, M., Murray, S.S., Jones, C., Forman, W., Van Speybroeck, L., 2005, ApJ, 628, 655
- Voit, M., 2005, Rev. Mod. Phys., 77, 207
- White, D.A., Jones, C. & Forman W., 1997, MNRAS, 292, 419
- White, D.A., 2000, MNRAS, 312, 663
- Zhang, Y.Y., Böhringer, H., Finoguenov, A. et al. 2006a, A&A, submitted, [astro-ph/0603275]
- Zhang, Y.Y., Finoguenov, A., Böhringer, H. et al. 2006b, preprint